

SPECIAL LAGRANGIAN SUBMANIFOLDS OF THE NEARLY KAEHLER 6-SPHERE

LUC VRANCKEN

ABSTRACT. In this paper, we study Lagrangian submanifolds of the nearly Kähler 6-sphere $S^6(1)$. It is well known that such submanifolds, which are 3-dimensional, are always minimal and admit a symmetric cubic form. Following an idea of Bryant, developed in the study of Lagrangian submanifolds of \mathbb{C}^3 , we then investigate those Lagrangian submanifolds which at each point admit an isometry preserving this cubic form. We obtain that all such Lagrangian submanifolds can be obtained starting from complex curves in $S^6(1)$ or from holomorphic curves in $\mathbb{C}P^2(4)$. As a corollary we classify the Lagrangian submanifolds which admit a Sasakian structure which is compatible with the induced metric. This last result generalizes theorems obtained by Deshmukh and ElHadi.

Subject class: 53B25, 53D12

1. INTRODUCTION

It is well known that starting from the Cayley numbers, it is possible to introduce an almost complex structure J on the 6-dimensional sphere $S^6(1)$ which is compatible with the standard metric. It was shown by Calabi and Gluck, see [5] that this structure, from a geometric viewpoint, is the best possible complex structure on $S^6(1)$. In the study of submanifolds, it is then natural to study submanifolds for which J preserves the tangent space (and hence also the normal space) and those for which J interchanges the tangent and normal spaces. The first class are called almost complex submanifolds and it was shown by Gray that they have to be two dimensional (complex one dimensional). Further results about these complex curves were obtained amongst others in [4], [10] and [2].

The second class of submanifolds mentioned, which by its definition have to be 3-dimensional, are called Lagrangian submanifolds. They were first investigated by Ejiri, [13], who showed that a Lagrangian submanifold is always orientable and minimal. Moreover, as is the case for Lagrangian submanifolds of complex space forms, we have that the 3-form C defined by

$$C(X, Y, Z) = \langle h(X, Y), JZ \rangle,$$

where h denotes the second fundamental form of the immersion is always symmetric. This implies that at every point p of M , we can introduce a symmetric polynomial f_p by

$$f_p(x, y, z) = \langle h(xe_1 + ye_2 + ze_3, xe_1 + ye_2 + ze_3), J(xe_1 + ye_2 + ze_3) \rangle,$$

where $\{e_1, e_2, e_3\}$ is an orthonormal basis at the point p . As M is minimal we see that the trace of this polynomial with respect to the metric vanishes. As far as such symmetric polynomials with vanishing trace on a 3-dimensional real vector space are concerned, we quote the following result by Bryant [3]:

Theorem 1. *Let $p \in M$ and assume that there exist an orientation preserving isometry which preserves f_p . Then there exists an orthonormal basis of $T_p M$ such that either*

- (i) $f_p = 0$, in this case f_p is preserved by every isometry,
- (ii) $f_p = \lambda(2x^3 - 3xy^2 - 3xz^2)$, for some positive number λ in which case f_p is preserved by a 1-parameter group of rotations,
- (iii) $f_p = 6\lambda xyz$ for some positive number λ , in which case f_p is preserved by the discrete group A_4 of order 12,
- (iv) $f_p = \lambda(x^3 - 3xy^2)$ for some positive number λ , in which case f_p is preserved by the discrete group S_3 of order 6,

Mathematisch Instituut, Universiteit Utrecht, Budapestlaan 6, 3584CD Utrecht, The Netherlands, Email: vrancken@math.uu.nl.

- (v) $f_p = \lambda(2x^3 - 3xy^2 - 3xz^2) + \mu(y^3 - 3xy^2)$ for some $\lambda, \mu > 0$, with $\mu \neq \sqrt{2}\lambda$, in which case f_p is preserved by the group \mathbb{Z}_3 ,
- (vi) $f_p = \lambda(2x^3 - 3xy^2 - 3xz^2) + 6\mu xyz$, for some $\lambda, \mu > 0$, with $\lambda \neq \mu$, in which case f_p is preserved by the group \mathbb{Z}_2 of order 2.

In this paper, we will assume that one of the special cases of the above theorem is satisfied at every point of the Lagrangian submanifold. We call M a Lagrangian submanifold of Type (k) if and only if Theorem 1(k) is satisfied at every point p of M . As it turns out, several of these classes of Lagrangian submanifolds have been previously studied. For example, the Lagrangian submanifolds of Type (iv) correspond to the Lagrangian submanifolds which satisfy Chen's equality which were previously studied in [11], [7] and [8].

The paper is organized as follows. In Section 2, we recall the construction of the almost complex structure on $S^6(1)$, starting from the Cayley multiplication, as well as some basic facts about Lagrangian submanifolds. Next, we start our investigation of Lagrangian submanifolds of Type (k). We start in Section 3 with collecting all known results translated to this setting. Next we show that, in contrast to the \mathbb{C}^3 -case studied by Bryant [3], there does not exist any Lagrangian submanifold of Type (v). As a corollary we remark that every Lagrangian submanifold of Type (k) which has constant scalar curvature must be equivariant and thus congruent to one of the 5 immersions of $SU(2)$ into $S^6(1)$, first described by Mashimo in [16]. This gives evidence to the following conjecture:

Conjecture 1. *Let $\psi : M^3 \rightarrow S^6(1)$ be a Lagrangian immersion with constant scalar curvature. Then ψ is congruent with an open part of one of the 5 previously mentioned equivariant immersions.*

The above conjecture can be seen as the analog for Lagrangian submanifolds of the well-known conjecture by Chern which states that the set of all possible values for the scalar curvature of a compact minimal hypersurface in a sphere is a discrete set.

As a Lagrangian submanifold M of $S^6(1)$ is always 3-dimensional, and thus odd-dimensional, it is a natural question to ask whether M admits a Sasakian structure compatible with the induced metric. This problem was first considered in [12], where some partial results were obtained. In Section 6, we completely classify those Lagrangian submanifolds of $S^6(1)$ which admit such a Sasakian structure.

2. PRELIMINARIES

We give a brief exposition of how the standard nearly Kähler structure on $S^6(1)$ arises in a natural manner from the Cayley multiplication. We also describe how we can use the vector cross product on \mathbb{R}^7 in order to define the Sasakian structure on $S^5(1)$. For further details about the Cayley numbers and their automorphism group G_2 , we refer the reader to [18] and [15].

The multiplication on the Cayley numbers \mathcal{O} may be used to define a vector cross product \times on the purely imaginary Cayley numbers \mathbb{R}^7 using the formula

$$(1) \quad u \times v = \frac{1}{2}(uv - vu),$$

while the standard inner product on \mathbb{R}^7 is given by

$$(2) \quad (u, v) = -\frac{1}{2}(uv + vu).$$

It is now elementary [15] to show that

$$(3) \quad u \times (v \times w) + (u \times v) \times w = 2(u, w)v - (u, v)w - (w, v)u,$$

and that the triple scalar product $(u \times v, w)$ is skew symmetric in u, v, w .

Conversely, Cayley multiplication on \mathcal{O} is given in terms of the vector cross product and the inner product by

$$(4) \quad (r + u)(s + v) = rs - (u, v) + rv + su + (u \times v), \quad r, s \in \text{Re}(\mathcal{O}), u, v \in \text{Im}(\mathcal{O})$$

In view of (1), (2) and (4), it is clear that the group G_2 of automorphisms of \mathcal{O} is precisely the group of isometries of \mathbb{R}^7 preserving the vector cross product.

An ordered basis u_1, \dots, u_7 is said to be a G_2 -frame if

$$(5) \quad u_3 = u_1 \times u_2, \quad u_5 = u_1 \times u_4, \quad u_6 = u_2 \times u_4, \quad u_7 = u_3 \times u_4.$$

For example, the standard basis e_1, \dots, e_7 of \mathbb{R}^7 is a G_2 -frame. Moreover, if u_1, u_2, u_4 are mutually orthogonal unit vectors with u_4 orthogonal to $u_1 \times u_2$, then u_1, u_2, u_4 determine a unique G_2 -frame u_1, \dots, u_7 and (\mathbb{R}^7, \times) is generated by u_1, u_2, u_4 subject to the relations :

$$(6) \quad u_i \times (u_j \times u_k) + (u_i \times u_j) \times u_k = 2\delta_{ik}u_j - \delta_{ij}u_k - \delta_{jk}u_i.$$

Therefore, for any G_2 -frame, we have the following very usefull multiplication table [18] :

x	u_1	u_2	u_3	u_4	u_5	u_6	u_7
u_1	0	u_3	$-u_2$	u_5	$-u_4$	$-u_7$	u_6
u_2	$-u_3$	0	u_1	u_6	u_7	$-u_4$	$-u_5$
u_3	u_2	$-u_1$	0	u_7	$-u_6$	u_5	$-u_4$
u_4	$-u_5$	$-u_6$	$-u_7$	0	u_1	u_2	u_3
u_5	u_4	$-u_7$	u_6	$-u_1$	0	$-u_3$	u_2
u_6	u_7	u_4	$-u_5$	$-u_2$	u_3	0	$-u_1$
u_7	$-u_6$	u_5	u_4	$-u_3$	$-u_2$	u_1	0

The standard nearly Kähler structure on $S^6(1)$ is then obtained as follows :

$$Ju = x \times u, \quad u \in T_x S^6(1), \quad x \in S^6(1).$$

It is clear that J is an orthogonal almost complex structure on $S^6(1)$. In fact J is a nearly Kähler structure in the sense that the $(2, 1)$ -tensor field G on $S^6(1)$ defined by

$$G(X, Y) = (\tilde{\nabla}_X J)Y,$$

where $\tilde{\nabla}$ is the Levi-Civita connection on $S^6(1)$ is skew-symmetric. A straightforward computation also shows that

$$G(X, Y) = X \times Y - \langle x \times X, Y \rangle x.$$

For more information on the properties of J , J and G , we refer to [2] and [10].

Let M be a submanifold of $S^6(1)$. Then, M is called Lagrangian provided that J interchanges at every point p of M the tangent and the normal space. It is immediately clear that a Lagrangian submanifold is 3-dimensional. It was also shown by Ejiri, [13] that M is minimal, orientable and that for tangent vector fields X and Y to M , $G(X, Y)$ is normal to M . Decomposing $\tilde{\nabla}_X Y$ and $\tilde{\nabla}_X JY$ into a tangential and a normal component it follows that

$$\nabla_X^\perp JY = J\nabla_X Y + G(X, Y),$$

and that the form defined by

$$C(X, Y, Z) = \langle h(X, Y), JZ \rangle,$$

is symmetric in X, Y and Z .

Now, we finish this section, by recalling some basic facts about the Hopf lift, which we will need in some of the later sections. It is well-known (see for instance [1], page 32 or [11]) that the complex structure of \mathbb{C}^3 induces a Sasakian structure (φ, ξ, η, g) on $S^5(1)$ starting from \mathbb{C}^3 . This structure can also be expressed using the vector cross product. We consider $S^5(1)$ as the hypersphere in $S^6(1) \subset \mathbb{R}^7$ given by the equation $x_4 = 0$ and define :

$$j : S^5(1) \rightarrow \mathbb{C}^3 : (x_1, x_2, x_3, 0, x_5, x_6, x_7) \mapsto (x_1 + ix_5, x_2 + ix_6, x_3 + ix_7).$$

Then at a point $p = (x_1, x_2, x_3, 0, x_5, x_6, x_7)$, the structure vector field ξ is given by

$$\xi(p) = (x_5, x_6, x_7, 0, -x_1, -x_2, -x_3) = e_4 \times p,$$

and for any tangent vector v , we get that

$$\varphi(v) = v \times e_4 - \langle v \times e_4, p \rangle p.$$

Following [19], we call a submanifold M^n of $S^5(1)$ invariant if $\varphi(T_p M) \subset T_p M$ for every p . If n is odd, then ξ is automatically tangent to M . Assume $n = 3$. The Hopf fibration $h : S^5(1) \rightarrow \mathbb{C}P^2(4)$ annihilates ξ , i.e. $dh(\xi) = 0$. Then if M^3 is invariant, $h(M^3)$ is a holomorphic curve. Conversely, let

$\phi : N_1 \rightarrow \mathbb{C}P^2(4)$ be a holomorphic curve, let PN_1 be the circle bundle over N_1 induced by the Hopf fibration and let ϕ be the immersion such that the following diagram commutes :

$$\begin{array}{ccc} PN_1 & \xrightarrow{\psi} & S^5(1) \\ \downarrow & & \downarrow h \\ N_1 & \xrightarrow{\phi} & \mathbb{C}P^2(4) \end{array}$$

Then ψ is an invariant immersion in the Sasakian space form $S^5(1)$ with structure vector field ξ tangent along ξ .

3. LAGRANGIAN SUBMANIFOLDS OF TYPE (i) TO (v)

In this section, we assume that M is a Lagrangian submanifold of $S^6(1)$ of Type (i) to (v). First, we assume that M is either Type (i) or Type (iii). This means that at each point p there exists an orthonormal basis $\{e_1, e_2, e_3\}$ such that

$$\begin{aligned} h(e_1, e_2) &= \lambda J e_3, & h(e_2, e_3) &= \lambda J e_1, & h(e_3, e_1) &= \lambda J e_2, \\ h(e_1, e_1) &= 0, & h(e_2, e_2) &= 0, & h(e_3, e_3) &= 0. \end{aligned}$$

Substituting this in the Gauss equation implies that

$$\widehat{R}(X, Y)Z = (1 - \lambda^2)(\langle Y, Z \rangle X - \langle X, Z \rangle Y).$$

Consequently, Schur's lemma implies that λ is a constant and that M has constant sectional curvature. Thus, from the classification of Lagrangian submanifolds with constant sectional curvature, see [13], it follows that either M is totally geodesic, corresponding to the case that $\lambda = 0$, or M is isometric to an open part of the 3-dimensional sphere with constant sectional curvature $\frac{1}{16}$, corresponding to the case that $\lambda = \sqrt{\frac{15}{16}}$. Moreover, in both cases, the immersion is unique up to applying an isometry of G_2 . Using the description of [14] for the immersion with constant sectional curvature $\frac{1}{16}$, we can summarize the above as follows:

Theorem 2. *Let $\psi : M \rightarrow S^6(1)$ be a Lagrangian immersion of Type (i). Then M is congruent to an open part of the immersion $\psi_1 : S^3(1) \rightarrow S^6(1) : (y_1, y_2, y_3, y_4) \mapsto (y_1, 0, y_2, 0, y_3, 0, y_4)$.*

Theorem 3. *Let $\psi : M \rightarrow S^6(1)$ be a Lagrangian immersion of Type (i). Denote by $\phi : S^2 \rightarrow S^6(1)$ the Veronese immersion of S^2 into $S^6(1)$ with constant Gaussian curvature $\frac{1}{6}$ and immersed such that J preserves the tangent space. Let α denote the second fundamental form of this immersion. Then,*

$$\psi_2 : US^2 \rightarrow S^6(\frac{1}{6}) \cos \gamma \phi + \sin \gamma v \times \frac{\alpha(v, v)}{\|\alpha(v, v)\|},$$

where $\cos^2 \gamma = \frac{5}{9}$ is a Lagrangian submanifold with constant sectional curvature $\frac{1}{16}$. Moreover, the immersion ψ is locally congruent to an open part of the immersion ψ_2 .

Next, we consider the case that M is a Lagrangian submanifold of Type (iv). This means that at each point p there exists an orthonormal basis $\{e_1, e_2, e_3\}$ such that

$$\begin{aligned} h(e_1, e_1) &= \lambda e_1, & h(e_1, e_2) &= -\lambda e_2, & h(e_3, e_1) &= 0, \\ h(e_2, e_2) &= -\lambda e_1, & h(e_2, e_3) &= 0, & h(e_3, e_3) &= 0, \end{aligned}$$

where λ is a positive number. From the above expression, it is clear that M realizes at every point the equality in the following inequality, which was obtained by Chen in [6] and which specialized to 3-dimensional submanifolds of $S^6(1)$, states

$$(7) \quad \delta_M(p) \leq \frac{9}{4} H^2(p) + 2,$$

for each $p \in M$, where H denotes the length of the mean curvature vector and δ_M is the Riemannian invariant, introduced by Chen in [6], defined by

$$\delta_M(p) = \tau(p) - (\inf K)(p).$$

Here

$$(\inf K)(p) = \inf \{K(\pi) \mid \pi \text{ is a 2-dimensional subspace of } T_p M\},$$

where $K(\pi)$ is the sectional curvature of π , and $\tau(p) = \sum_{i < j} K(e_i \wedge e_j)$ denotes the scalar curvature defined in terms of an orthonormal basis $\{e_1, e_2, e_3\}$ of the tangent space $T_p M$. Submanifolds realizing the equality are called submanifolds satisfying Chen's equality. As a Lagrangian submanifold of $S^6(1)$ is always minimal, it is clear that M realizes Chen's equality if and only if $\delta_M = 2$. Such Lagrangian submanifolds were classified in [11].

Theorem 4. *Let $\phi : N_1 \rightarrow \mathbb{C}P^2(4)$ be a holomorphic curve in $\mathbb{C}P^2(4)$. Let PN_1 be the circle bundle over N_1 induced by the Hopf fibration $p : S^5(1) \rightarrow \mathbb{C}P^2(4)$ and let ψ be the isometric immersion such that the following diagram commutes:*

$$\begin{array}{ccc} PN_1 & \xrightarrow{\psi} & S^5(1) \\ \downarrow & & \downarrow p \\ N_1 & \xrightarrow{\phi} & \mathbb{C}P^2(4) \end{array}$$

Then, there exists a totally geodesic embedding i of $S^5(1)$ into the nearly Kähler 6-sphere such that the immersion $i \circ \psi : PN_1 \rightarrow S^6(1)$ is a 3-dimensional Lagrangian immersion in $S^6(1)$ satisfying Chen's equality. Moreover, every Lagrangian submanifold which is contained in a totally geodesic $S^5(1)$ in $S^6(1)$ can be obtained in this way.

Theorem 5. *Let $\bar{\phi} : N_2 \rightarrow S^6(1)$ be an almost complex curve (with second fundamental form α) without totally geodesic points. Denote by UN_2 the unit tangent bundle over N_2 and define a map*

$$\bar{\psi} : UN_2 \rightarrow S^6(1) \times \frac{\alpha(v,v)}{\|\alpha(v,v)\|}.$$

Then $\bar{\psi}$ is a Lagrangian immersion into $S^6(1)$ satisfying Chen's equality. Moreover, the immersion is linearly full in $S^6(1)$. Conversely, every Lagrangian immersion satisfying Chen's equality can be obtained in this way.

Using the above theorems it is straightforward to compute that a Lagrangian submanifold of Type (iv) has constant scalar curvature if the underlying complex curve in $\mathbb{C}P^2(4)$ or $S^6(1)$, depending on the case which we are considering, is not totally geodesic and has constant Gaussian curvature.

Finally, we consider the case that M is a Lagrangian submanifold of Type (ii) or Type (v). In that case, we know that at each point p of M there exists an orthonormal basis $\{e_1, e_2, e_3\}$ such that

$$\begin{aligned} h(e_1, e_1) &= 2\lambda J e_1, & h(e_1, e_2) &= -\lambda J e_1, & h(e_3, e_1) &= -\lambda J e_3, \\ h(e_2, e_2) &= -\lambda J e_1 + \mu e_2, & h(e_2, e_3) &= -\mu J e_3, & h(e_3, e_3) &= -\lambda J e_1 - \mu J e_2, \end{aligned}$$

where λ is a positive number and $\mu \neq \sqrt{2}\lambda$. If M is of Type (ii) then $\mu = 0$, whereas if M is of Type (vi) then $\mu > 0$. Then, a straightforward computation using the Gauss equation shows that:

$$\begin{aligned} \text{Ric}(e_1, e_1) &= 1 - 3\lambda^2, & \text{Ric}(e_1, e_2) &= 0, & \text{Ric}(e_3, e_1) &= 0, \\ \text{Ric}(e_2, e_2) &= 1 - \lambda^2 - \mu^2, & \text{Ric}(e_2, e_3) &= 0, & \text{Ric}(e_3, e_3) &= 1 - \lambda^2 - \mu^2, \end{aligned}$$

where the Ricci tensor Ric is defined by

$$\text{Ric}(Y, Z) = \frac{1}{2} \text{trace}\{X \mapsto R(X, Y)Z\}$$

As $\mu^2 \neq 2\lambda^2$, we see as a consequence that the 1-1 symmetric tensor field P associated with the Ricci tensor has at each point two different eigenvalues, one with multiplicity 1, the other with multiplicity 2. In particular this means that M is a quasi Einstein Lagrangian manifold. As, in this case, M does not satisfy Chen's equality it follows from the main theorem of [9] that M can be obtained as follows:

Theorem 6. *Let $\phi : N^2 \rightarrow S^6$ be a superminimal almost complex curve in $S^6(1)$ without totally geodesic points and different from the Veronese immersion. Denote by UN^2 the unit tangent bundle of N^2 . Define*

$$\psi_\gamma : UN^2 \rightarrow S^6 := \cos \gamma \phi + \sin \gamma v \times \frac{\alpha(v,v)}{\|\alpha(v,v)\|},$$

where $\cos^2 \gamma = \frac{5}{9}$. Then ψ is a Lagrangian immersion on an open dense subset of UN^2 . Moreover ψ is a Lagrangian immersion of Type (iv). Conversely, every Lagrangian submanifold of Type (iv) can be obtained in this way.

Theorem 7. Let $\psi : M \rightarrow S^6(1)$ be a Lagrangian immersion of Type (ii). Denote by S^3 the 3-dimensional sphere and consider the map into $S^6(1)$ defined by:

$$\tilde{\psi} : S^3 \rightarrow S^6(1) : (y_1, y_2, y_3, y_4) \mapsto (x_1, x_2, x_3, x_4, x_5, x_6, x_7),$$

where

$$\begin{aligned} x_1 &= \frac{1}{9}(5y_1^2 + 5y_2^2 - 5y_3^2 - 5y_4^2 + 4y_1), \\ x_2 &= -\frac{2}{3}y_2, \\ x_3 &= \frac{2\sqrt{5}}{9}(y_1^2 + y_2^2 - y_3^2 - y_4^2 - y_1), \\ x_4 &= \frac{\sqrt{3}}{9\sqrt{2}}(-10y_3y_1 - 2y_3 - 10y_2y_4), \\ x_5 &= \frac{\sqrt{3}\sqrt{5}}{9\sqrt{2}}(2y_1y_4 - 2y_4 - 2y_2y_3), \\ x_6 &= \frac{\sqrt{3}\sqrt{5}}{9\sqrt{2}}(2y_1y_3 - 2y_3 + 2y_2y_4), \\ x_7 &= \frac{\sqrt{3}}{9\sqrt{2}}(10y_1y_4 + 2y_4 - 10y_2y_3), \end{aligned}$$

and $y_1^2 + y_2^2 + y_3^2 + y_4^2 = 1$. Then $\tilde{\psi}$ defines a Lagrangian immersion of Type (ii) with constant scalar curvature. Conversely every Lagrangian immersion of Type (ii) is congruent with an open part of $\tilde{\psi}$.

Note that the induced metric on S^3 by the above immersion is not the standard metric on S^3 .

4. LAGRANGIAN SUBMANIFOLDS OF TYPE (vi)

In this section, we assume that M is a Lagrangian submanifold of Type (vi). It is easy to see that this implies that at each point p there exists an orthonormal basis $\{e_1, e_2, e_3\}$ such that

$$\begin{aligned} h(e_1, e_1) &= \lambda_1 J e_1, & h(e_1, e_2) &= \lambda_2 J e_2, & h(e_3, e_1) &= \lambda_3 J e_3, \\ h(e_2, e_2) &= \lambda_2 J e_1, & h(e_2, e_3) &= 0, & h(e_3, e_3) &= \lambda_3 J e_1, \end{aligned}$$

where $0 < \lambda_1 = -\lambda_2 - \lambda_3$ and $0 \neq \lambda_2 \neq \lambda_3 \neq 0$.

By a straightforward computation we obtain the following:

Lemma 1. Let $\{e_1, e_2, e_3\}$ be the orthonormal basis defined previously. Then it follows that

$$\begin{aligned} \text{Ric}(e_1, e_1) &= 1 - \lambda_2^2 - \lambda_3^2 - \lambda_2\lambda_3, & \text{Ric}(e_1, e_2) &= 0, & \text{Ric}(e_3, e_1) &= 0, \\ \text{Ric}(e_2, e_2) &= 1 - \lambda_2^2, & \text{Ric}(e_2, e_3) &= 0, & \text{Ric}(e_3, e_3) &= 1 - \lambda_3^2. \end{aligned}$$

Using the various conditions on λ_2 and λ_3 , we see that the 1-1 symmetric tensor field P associated with the Ricci tensor has at each point three different eigenvalues, all with multiplicity 1. Hence there exist orthonormal vector fields $\{E_1, E_2, E_3\}$ defined on a neighborhood of the point p and a non vanishing differentiable functions λ_2, λ_3 , with $\lambda_2 + \lambda_3 < 0$ and $\lambda_2 - \lambda_3 \neq 0$ such that

$$\begin{aligned} h(E_1, E_1) &= -(\lambda_2 + \lambda_3)J E_1, & h(E_1, E_2) &= \lambda_2 J E_2, & h(E_3, E_1) &= \lambda_3 J E_3, \\ h(E_2, E_2) &= \lambda_2 J E_1, & h(E_2, E_3) &= 0, & h(E_3, E_3) &= \lambda_3 J E_1. \end{aligned}$$

As $G(E_1, E_2) = \pm J E_3$, we may, by replacing E_3 with $-E_3$ if necessary, assume that $G(E_1, E_2) = J E_3$. We then introduce local functions a_1, \dots, c_3 by

$$\begin{aligned} \nabla_{E_1} E_1 &= a_1 E_2 + a_2 E_3, & \nabla_{E_1} E_2 &= -a_1 E_1 + a_3 E_3, & \nabla_{E_1} E_3 &= -a_2 E_1 - a_3 E_2, \\ \nabla_{E_2} E_1 &= b_1 E_2 + b_2 E_3, & \nabla_{E_2} E_2 &= -b_1 E_1 + b_3 E_3, & \nabla_{E_2} E_3 &= -b_2 E_1 - b_3 E_2, \\ \nabla_{E_3} E_1 &= c_1 E_2 + c_2 E_3, & \nabla_{E_3} E_2 &= -c_1 E_1 + c_3 E_3, & \nabla_{E_3} E_3 &= -c_2 E_1 - c_3 E_2, \end{aligned}$$

Computing now all components of the Gauss equation, it then follows by a long but straightforward computation that the functions $\lambda_2, \lambda_3, a_1, \dots, c_3$ have to satisfy the following system of partial differential equations:

$$(8) \quad E_2(a_1) - E_1(b_1) = 1 - 2\lambda_2^2 - \lambda_2\lambda_3 + a_1^2 + b_1^2 + b_2c_1 - b_2a_3 + a_2b_3 - a_3c_1,$$

$$(9) \quad E_3(a_2) - E_1(c_2) = 1 - 2\lambda_3^2 - \lambda_2\lambda_3 + a_2^2 + c_2^2 + b_2c_1 + b_2a_3 - a_1c_3 + a_3c_1,$$

$$(10) \quad E_2(a_2) - E_1(b_2) = b_1a_3 + b_1b_2 + a_1a_2 - a_1b_3 + c_2b_2 - c_2a_3,$$

$$(11) \quad E_3(a_1) - E_1(c_1) = a_2c_3 - a_3c_2 + a_1a_2 + b_1c_1 + c_1c_2 + b_1a_3,$$

$$(12) \quad E_2(c_2) - E_3(b_2) = b_1c_3 - b_3c_1 - a_2b_2 + a_2c_1 - b_2b_3 - c_2c_3,$$

$$(13) \quad E_2(c_1) - E_3(b_1) = b_3c_2 - c_3b_2 + a_1c_1 - a_1b_2 - c_1c_3 - b_1b_3,$$

$$(14) \quad E_3(a_3) - E_1(c_3) = a_1c_2 - a_2c_1 + a_2a_3 + a_3b_3 + b_3c_1 + c_2c_3,$$

$$(15) \quad E_1(b_3) - E_2(a_3) = b_1a_2 - a_1b_2 - a_1a_3 - b_1b_3 + a_3c_3 - c_3b_2,$$

$$(16) \quad E_3(b_3) - E_2(c_3) = 1 + \lambda_2\lambda_3 + b_3^2 + c_3^2 + a_3b_2 - a_3c_1 + b_1c_2 - b_2c_1.$$

The number of unknowns in the above equations can be reduced using the Codazzi equation which states that $(\nabla h)(X, Y, Z) = (\nabla_X h)(Y, Z) = \nabla_X^{\perp} h(Y, Z) - h(\nabla_X Y, Z) - h(Y, \nabla_X Z)$ is totally symmetric in X, Y and Z . In particular we obtain that

Lemma 2. *Let $\{E_1, E_2, E_3\}$ be the local orthonormal basis defined previously. Then, we have that there exists a function c such that*

$$b_2 = -c_1 = -a_3 = \frac{1}{4},$$

$$a_2 = \left(1 - \frac{\lambda_3}{\lambda_2}\right)b_3,$$

$$a_1 = \left(\frac{\lambda_2}{\lambda_3} - 1\right)c_3,$$

$$c_2 = c\lambda_3,$$

$$b_1 = c\lambda_2.$$

Moreover, the functions λ_2 and λ_3 satisfy the following system of differential equations:

$$E_1(\lambda_2) = -c\lambda_2(3\lambda_2 + \lambda_3),$$

$$E_2(\lambda_2) = 3c_3\lambda_2\left(\frac{\lambda_2}{\lambda_3} - 1\right),$$

$$E_3(\lambda_2) = b_3(\lambda_2 - \lambda_3),$$

$$E_1(\lambda_3) = -c\lambda_3(3\lambda_3 + \lambda_2),$$

$$E_2(\lambda_3) = c_3(\lambda_2 - \lambda_3),$$

$$E_3(\lambda_3) = 3b_3\lambda_3\left(1 - \frac{\lambda_3}{\lambda_2}\right).$$

Proof. As,

$$(\nabla_{E_2} h)(E_3, E_3) = E_2(\lambda_3)JE_1 + b_1\lambda_3JE_2 + (3b_2 - 1)\lambda_3JE_3,$$

and

$$(\nabla_{E_3} h)(E_2, E_3) = c_3(\lambda_2 - \lambda_3)JE_1 + c_2\lambda_2JE_2 + c_1\lambda_3JE_3,$$

it follows from the Codazzi equation $(\nabla_{E_2} h)(E_3, E_3) = (\nabla_{E_3} h)(E_2, E_3)$ that

$$(17) \quad c_1 = 3b_2 - 1,$$

$$(18) \quad \lambda_3b_1 = c_2\lambda_2,$$

$$(19) \quad E_2(\lambda_3) = b_3(\lambda_2 - \lambda_3).$$

Similarly, we obtain from $(\nabla_{E_2} h)(E_3, E_2) = (\nabla_{E_3} h)(E_2, E_2)$ that

$$(20) \quad b_2 = 3c_1 + 1,$$

$$(21) \quad E_3(\lambda_2) = b_3(\lambda_2 - \lambda_3).$$

Combining (17) and (20) it then follows that $c_1 = -\frac{1}{4}$ and $b_2 = \frac{1}{4}$. The remaining equations follow similarly from the other Codazzi equations. \square

Using the previous lemma, the differential equations given by (10) to (16) now imply that

$$\begin{aligned} E_2(c) &= \frac{1}{2}b_3\left(\frac{1}{\lambda_2} - \frac{1}{\lambda_3}\right), \\ E_3(c) &= \frac{1}{2}c_3\left(\frac{1}{\lambda_3} - \frac{1}{\lambda_2}\right), \\ E_1(c_3) &= -cc_3\lambda_2 + \frac{1}{2}b_3, \\ E_1(b_3) &= -b_3c\lambda_3 - \frac{1}{2}c_3, \\ E_3(c_3) &= 3b_3c_3 - \frac{1}{2}c\frac{\lambda_2\lambda_3}{\lambda_2-\lambda_3}, \\ E_2(b_3) &= -3b_3c_3 + \frac{1}{2}c\frac{\lambda_2\lambda_3}{\lambda_2-\lambda_3}. \end{aligned}$$

We now compute some integrability conditions. As ∇ is torsion free, we know that for any function f , the following equations are satisfied:

$$\begin{aligned} 0 &= E_1(E_2(f)) - E_2(E_1(f)) - (\nabla_{E_1}E_2)(f) + (\nabla_{E_2}E_1)(f), \\ 0 &= E_1(E_3(f)) - E_3(E_1(f)) - (\nabla_{E_1}E_3)(f) + (\nabla_{E_3}E_1)(f), \\ 0 &= E_2(E_3(f)) - E_3(E_2(f)) - (\nabla_{E_2}E_3)(f) + (\nabla_{E_3}E_2)(f). \end{aligned}$$

It is straightforward to check that applying the above principle for the functions λ_2 and λ_3 does not yield any new equations. However, applying the first principle for the function b_3 yields the following differential equation:

$$\begin{aligned} \lambda_2\lambda_3(\lambda_2 - \lambda_3)(E_3(b_3) + E_2(c_3)) + \lambda_2^2\lambda_3^2E_1(c) \\ = (\lambda_2 - \lambda_3)(c_3^2\lambda_2(\lambda_2 - 4\lambda_3) + b_3^2\lambda_3(4\lambda_2 - \lambda_3)) \end{aligned}$$

Combining this equation, together with the remaining Gauss equations, we then obtain that

$$\begin{aligned} E_3(b_3) &= \frac{(-5-32b_3^2+16c_3^2)\lambda_3+2\lambda_2(5+16b_3^2-8c_3^2-8\lambda_3^2-8c^2\lambda_3^2)}{16(\lambda_2-\lambda_3)}, \\ E_2(c_3) &= \frac{(-5-32c_3^2+16b_3^2)\lambda_2+2\lambda_3(5+16c_3^2-8b_3^2-8\lambda_2^2-8c^2\lambda_2^2)}{16(\lambda_2-\lambda_3)}, \\ E_1(c) &= \frac{-5-32b_3^2+16c_3^2}{16\lambda_2} + \lambda_2\left(1 + c^2 + \frac{c^2}{\lambda_3^2}\right) \\ &\quad - \frac{-5-32c_3^2+16b_3^2}{16\lambda_3} + \lambda_3\left(1 + c^2 + \frac{b_3^2}{\lambda_2^2}\right). \end{aligned}$$

Checking now the integrability conditions for c , c_3 and b_3 it immediately follows that $c = c_3 = b_3 = 0$. Substituting these values in to the Gauss equations it follows that $\lambda_2 = -\lambda_3$ which is a contradiction. Therefore, we have shown the following theorem:

Theorem 8. *There does not exist a Lagrangian submanifold of Type (vi) in $S^6(1)$.*

5. LAGRANGIAN SUBMANIFOLDS ADMITTING A SASAKIAN STRUCTURE

Throughout this section, we will assume that M is a Lagrangian submanifold of $S^6(1)$ which admit a Sasakian structure which is compatible with the induced metric $\langle \cdot, \cdot \rangle$ on M . This implies, see [1], that there exists a unit-length vector field ξ on M , a 1-form η and an endomorphism Φ satisfying:

$$\begin{aligned} \eta(X) &= \langle X, \xi \rangle, \\ \Phi^2 &= -I + \eta \otimes \xi, \\ \Phi(\xi) &= 0, \\ \eta \circ \Phi &= 0, \\ \langle \Phi X, \Phi Y \rangle &= \langle X, Y \rangle - \eta(X)\eta(Y), \\ (\nabla_X \phi)Y &= \eta(Y)X - \langle X, Y \rangle \xi. \end{aligned}$$

Moreover, it is well known, see [1], that the above equations imply that

$$(22) \quad \nabla_X \xi = -\Phi X,$$

$$(23) \quad R(X, Y)\xi = \langle \xi, Y \rangle X - \langle X, \xi \rangle Y.$$

From [1], we recall the following theorem that the previous equation together with the fact that ξ is a unit-length Killing vector field are the principal criteria for determining whether an odd-dimensional manifold admits a Sasakian structure compatible with a given metric.

Lemma 3. *Let M^3 be a Riemannian manifold admitting a unit length Killing vector field ξ such that*

$$R(X, Y)\xi = \langle \xi, Y \rangle X - \langle \xi, X \rangle Y,$$

then M admits a Sasakian structure which is compatible with the given metric.

Taking now an orthonormal basis $\{e_1, e_2, e_3\}$ at a point p of M^3 such that $e_3 = \xi$, it immediately follows from (23) that

$$\begin{aligned} \text{Ric}(e_1, e_1) &= \frac{1}{2}(\langle R(e_1, e_2)e_2, e_1 \rangle + 1), & \text{Ric}(e_1, e_2) &= 0, & \text{Ric}(e_3, e_1) &= 0, \\ \text{Ric}(e_2, e_2) &= \frac{1}{2}(\langle R(e_1, e_2)e_2, e_1 \rangle + 1), & \text{Ric}(e_2, e_3) &= 0, & \text{Ric}(e_3, e_3) &= 1. \end{aligned}$$

This implies that all sectional curvatures at the point p equal 1 or the associated endomorphism P has two eigenvalues: one with multiplicity 1 and one with multiplicity 2. Moreover, in the second case, the eigenvalue with multiplicity 1 equals 1 and the vector field ξ spans the corresponding 1-dimensional eigenspace. We now recall the following lemma from [9] about quasi-Einstein submanifolds.

Lemma 4. *Let M^3 be a 3-dimensional Lagrangian submanifold of S^6 with the second fundamental form h . Then the Ricci endomorphism P , associated with the Ricci tensor Ric , satisfies:*

- (i) *1 is an eigenvalue of P ,*
- (ii) *P has an eigenvalue with multiplicity at least 2,*

if and only if p is a totally geodesic point or there exist a orthonormal basis $\{e_1, e_2, e_3\}$ of $T_p M$ such that either

$$\begin{aligned} h(e_1, e_1) &= \lambda J e_1, & h(e_2, e_2) &= -\lambda J e_1, \\ h(e_1, e_2) &= -\lambda J e_2, & h(e_2, e_3) &= 0, \\ h(e_1, e_3) &= 0, & h(e_3, e_3) &= 0, \end{aligned}$$

where λ is a non-zero number. Moreover, in the second case, the 1-dimensional eigenspace is determined by e_3 .

It follows immediately from the above lemma that a Lagrangian submanifold which admits a Sasakian structure satisfies Chen's equality. Moreover, if necessary by restricting to an open dense subset, we may assume that either M is totally geodesic or in a neighborhood of any point p of M^3 there exist an orthonormal basis $\{E_1, E_2, E_3\}$ such that

$$\begin{aligned} h(E_1, E_1) &= \lambda J E_1, & h(E_2, E_2) &= -\lambda J E_1, & h(E_1, E_2) &= -\lambda J E_2, \\ h(E_2, E_3) &= 0, & h(E_1, E_3) &= 0, & h(E_3, E_3) &= 0, \end{aligned}$$

where $E_3 = \xi$. Now, we proceed as follows. We take the frame constructed in the previous lemma. As $G(E_3, E_1)$ is a normal vector which is orthogonal to both $J E_3$ and $J E_1$ it follows that $G(E_3, E_1) = \pm E_2$. Therefore, if necessary by changing the sign of E_2 , we may assume that $G(E_3, E_1) = J E_2$. It then follows that $G(E_1, E_2) = J E_3$ and $G(E_2, E_3) = J E_1$. Moreover, we also have that $\Phi E_1 = \epsilon E_2$, where $\epsilon = \pm 1$. It then follows from (22) that

$$\begin{aligned} \nabla_{E_1} E_3 &= -\epsilon E_2, \\ \nabla_{E_2} E_3 &= \epsilon E_1, \\ \nabla_{E_3} E_3 &= 0. \end{aligned}$$

Introducing now functions a, b and c , it follows that we can express the other components of the connection ∇ respectively by

$$\begin{aligned} \nabla_{E_1} E_1 &= c E_2, & \nabla_{E_1} E_2 &= -c E_1 + \epsilon E_3, \\ \nabla_{E_2} E_1 &= d E_2 - \epsilon E_3, & \nabla_{E_2} E_2 &= -d E_1, \\ \nabla_{E_3} E_1 &= f E_2, & \nabla_{E_3} E_2 &= -f E_1. \end{aligned}$$

Using now the Codazzi equation, see also Lemma 5.3 of [8], it follows that $f = -\frac{1}{3}(1 + \epsilon)$. We now consider two different cases. First, we assume that $\epsilon = -1$. In that case, we have that

$$\begin{aligned}\tilde{\nabla}_{E_1} J E_3 &= G(E_1, E_3) + J \tilde{\nabla}_{E_1} E_3 \\ &= -J E_2 + J E_2 + h(E_1, E_3) = 0, \\ \tilde{\nabla}_{E_2} J E_3 &= G(E_2, E_3) + J \tilde{\nabla}_{E_2} E_3 \\ &= J E_1 - J E_1 + h(E_2, E_3) = 0, \\ \tilde{\nabla}_{E_3} J E_3 &= J \tilde{\nabla}_{E_3} E_3 = 0.\end{aligned}$$

Hence, $J E_3$ is a constant vector along M which is tangent to $S^6(1)$. Consequently M lies in the totally geodesic $S^5(1)$ which is obtained as the intersection of $S^6(1)$ with the linear hyperplane orthogonal to $J E_3$. Using now the classification of Lagrangian submanifolds contained in a totally geodesic subspace, see Theorem 4, we obtain that M is locally congruent to the Hopf lift of a complex curve in $\mathbb{C}P^2(4)$ to $S^5(1)$ and $S^5(1)$ is immersed in $S^6(1)$ as described in Section 2. As the Hopf lift of a complex curve is a Sasakian manifold, the converse is obvious.

Finally, we consider the case that $\epsilon = 1$. In this case, we denote the immersion by F and proceed as in [11] to show that the map $J E_3 = F \times E_3$ defines an almost complex curve. Specializing the formulas there, to our case, i.e. using that the functions a and b defined in [11] are respectively $a = 0$ and $b = 1$, we get that:

$$\begin{aligned}D_{E_1}(F \times E_3) &= 2E_1 \times E_3, \\ D_{E_2}(F \times E_3) &= 2E_2 \times E_3, \\ D_{E_3}(F \times E_3) &= 0, \\ D_{E_1}(E_1 \times E_3) &= (cE_2 + \lambda F \times E_1 - F) \times E_3 - E_1 \times E_2 = cE_2 \times E_3 - \lambda E_2 - 2F \times E_3, \\ D_{E_2}(E_1 \times E_3) &= (dE_2 - \lambda F \times E_2) \times E_3 = dE_2 \times E_3 - \lambda E_1, \\ D_{E_1}(E_2 \times E_3) &= (-cE_1 - \lambda F \times E_2) \times E_3 = -cE_1 \times E_3 - \lambda E_1, \\ D_{E_2}(E_2 \times E_3) &= (-dE_1 - \lambda F \times E_1 - F) \times E_3 + E_2 \times E_1 = -dE_1 \times E_3 + \lambda E_2 - 2F \times E_3, \\ D_{E_1} E_1 &= cE_2 + \lambda F \times E_1 - F = cE_2 + \lambda E_2 \times E_3 - F, \\ D_{E_1} E_2 &= -cE_1 - \lambda F \times E_2 + E_3 = -cE_1 + \lambda E_1 \times E_3 + E_3, \\ D_{E_2} E_1 &= dE_2 - \lambda F \times E_2 - E_3 = dE_2 + \lambda E_1 \times E_3 - E_3, \\ D_{E_2} E_2 &= -dE_1 - \lambda F \times E_1 - F = -dE_1 - \lambda E_2 \times E_3 - F.\end{aligned}$$

The above formulas now imply immediately that the complex curve defined by $J E_3$ is superminimal. A representation of superminimal complex curves was obtained in [4]. It now follows from [11] that M is obtained as in Theorem 5, starting from a superminimal complex curve in $S^6(1)$.

In order to show that we can start with an arbitrary superminimal complex curve it is sufficient to check that the vector field $\frac{\partial}{\partial t}$, defined in the proof of Theorem 2 of [11], is a Killing vector field satisfying the conditions of Lemma 3, which can be verified by a straightforward computation. Therefore, we have shown the following theorem:

Theorem 9. *Let $\psi : M^3 \rightarrow S^6(1)$ be a Lagrangian immersion. Then M admits a Sasakian structure compatible with the induced metric if and only if locally M is congruent with*

- (i) a totally geodesic immersion,
- (ii) the inverse image under the Hopf map of a holomorphic curve in $\mathbb{C}P^2(4)$, as described in Theorem 4,
- (iii) the unit second normal bundle of a superminimal complex curve in $S^6(1)$, as described in Theorem 5.

REFERENCES

- [1] D. E. Blair, Contact Manifolds im Riemannian Geometry *Lecture Notes in Mathematics*, 509, Springer, Berlin, 1976.
- [2] J. Bolton, L. Vrancken and L. M. Woodward, On almost complex curves in the nearly Kaehler 6-sphere *Quart. J. Math. Oxford Ser (2)*, 45(1994): 407-427.
- [3] R. L. Bryant, Second order families of special Lagrangian 3-folds, preprint.

- [4] R. L. Bryant, Submanifolds and special structures on the octonions *J. Differential Geometry*, 17 (1982): 185–232.
- [5] E. Calabi and H. Gluck, What are the best almost complex structures on the 6-sphere? in *Differential Geometry: geometry in mathematical physics and related topics*, Amer. Math. Soc, 1993: 99–106.
- [6] B. Y. Chen, Some pinching and classification theorems for minimal submanifolds *Archiv. Math. (Basel)*, 60(1993):568–578.
- [7] B. Y. Chen, F. Dillen, L. Verstraelen and L. Vrancken, Two equivariant totally real immersions into the nearly Kähler 6-sphere and their characterization *Japanese J. Math. (N.S)*, 21(1995), 207–222.
- [8] B. Y. Chen, F. Dillen, L. Verstraelen and L. Vrancken, Characterizing a class of totally real submanifolds of $S^6(1)$ by their sectional curvatures *Tôhoku Math. J.*, 47(1995): 185–198.
- [9] R. Deszcz, F. Dillen, L. Verstraelen and L. Vrancken, Quasi-Einstein totally real submanifolds of the nearly Kähler 6-sphere *Tôhoku Math. J.*, 51(1999): 461–478.
- [10] F. Dillen, L. Verstraelen and L. Vrancken, On problems of U. Simon concerning minimal submanifolds of the nearly Kähler 6-sphere *Bull. Amer. Math. Soc.*, 19(1988): 433–438.
- [11] F. Dillen and L. Vrancken, Totally real submanifolds in $S^6(1)$ satisfying Chen's equality *Trans. Amer. Math. Soc.*, 348(1996): 1633–1646.
- [12] S. Deshmukh and K. El Hadi, A Sasakian structure on a 3-dimensional totally real submanifold of the nearly Kähler 6-sphere *Panamer. Math. J.*, 2(1992): 43–52.
- [13] N. Ejiri, Totally real submanifolds in a 6-sphere *Proc. Amer. Math. Soc.*, 83 (1981): 759–763.
- [14] N. Ejiri, Equivariant minimal immersions of S^2 into S^{2m} *Trans. Amer. Math. Soc.*, 297(1986): 105–124.
- [15] R. Harvey and H. B. Lawson, Calibrated Geometries *Acta Math.*, 148(1982): 47–157.
- [16] K. Mashimo, Homogeneous totally real submanifolds of $S^6(1)$ *Tsukuba J. math.*, 9 (1985): 185–202.
- [17] B. Palmer, Calibrations and Lagrangian submanifolds in the six sphere *Tôhoku Math. J.*, 50(1998): 303–315.
- [18] R. M. Wood, Framing the exceptional Lie group G_2 *Topology*, 15 (1976): 303–320.
- [19] K. Yano and S. Ishihara, Invariant submanifolds of an almost contact manifold *Kodai. Math. Sem. Rep.*, 21 (1969): 350–364.