

**Universiteit Utrecht**



*Department  
of Mathematics*

**Arnoldi and Jacobi-Davidson methods  
for generalized eigenvalue problems**  
 *$Ax = \lambda Bx$  with singular  $B$*

by

**Joost Rommes**

Preprint

nr. 1339

March, 2005

revised version

January, 2007

# Arnoldi and Jacobi-Davidson methods for generalized eigenvalue problems $Ax = \lambda Bx$ with singular $B$

Joost Rommes\*

March, 2005 (revised January, 2007)

## Abstract

In many physical situations, a few specific eigenvalues of a large sparse generalized eigenvalue problem  $Ax = \lambda Bx$  are needed. If exact linear solves with  $A - \sigma B$  are available, implicitly restarted Arnoldi with purification is a common approach for problems where  $B$  is positive semidefinite. In this paper, a new approach based on implicitly restarted Arnoldi will be presented that avoids most of the problems due to the singularity of  $B$ . Secondly, if exact solves are not available, Jacobi-Davidson QZ will be presented as a robust method to compute a few specific eigenvalues. Results are illustrated by numerical experiments.

## 1 Introduction

Large sparse generalized eigenvalue problems of the form

$$Ax = \lambda Bx, \quad x \neq 0, \quad (1.1)$$

with  $A, B \in \mathbb{R}^{n \times n}$ ,  $x \in \mathbb{C}^n$  and  $\lambda \in \mathbb{C}$ , arise in physical situations like stability analysis of the discretized Navier-Stokes equations. Typically, the matrix  $A$  is nonsymmetric and of full rank, and  $B$  is singular. The pencil  $(A, B)$  is regular, that is,  $A - \gamma B$  is singular only for a finite number of  $\gamma \in \mathbb{C}$ . Because  $B$  is singular, (1.1) can have eigenvalues at infinity, which are of no physical relevance, but may lead to numerical difficulties. In practice, one is often interested in the few left- or rightmost *finite* eigenvalues that determine the stability, and hence one wants to avoid approximations to eigenvalues at infinity. This paper is concerned with the computation of a few left- or rightmost eigenvalues of large generalized eigenvalue problems.

One way to compute a few eigenvalues of (1.1) close to  $\sigma \in \mathbb{C}$  is to apply Arnoldi's method to the shift-and-invert transformation  $S = (A - \sigma B)^{-1}B$ :

$$Sx = \tilde{\lambda}x, \quad x \neq 0. \quad (1.2)$$

An eigenpair  $(\lambda, x)$  of (1.1) corresponds to an eigenpair  $(\tilde{\lambda} = (\lambda - \sigma)^{-1}, x)$  of (1.2). Hence, the infinite eigenvalues of (1.1) correspond to eigenvalues  $\tilde{\lambda} = 0$  of (1.2). Arnoldi's method may compute approximations  $\tilde{\theta}$  to  $\tilde{\lambda} = 0$ . These approximations are known as *spurious* eigenvalues and after back transformation via  $\theta = \tilde{\theta}^{-1} + \sigma$ , they may be hard to distinguish from wanted eigenvalues, which typically reside in the exterior of the spectrum. This problem has been addressed for the symmetric nondefective problem [5, 16] and for the defective problem [5]. The ideas presented there are extended to the nonsymmetric defective case in [14], where the implicitly restarted Arnoldi method [11, 22] is implemented with a  $B$  semi-inner product and purification. Purification is a technique to remove unwanted components from Arnoldi vectors and approximate eigenvectors, and will be explained in more detail in section 3. A new strategy will be presented that, by exploiting the structure of (1.1), reduces the corruption by unwanted components significantly.

---

\*Mathematical Institute, rommes@math.uu.nl

The scheme based on the Arnoldi method fails to be applicable if the linear system solves with  $A - \sigma B$ , e.g. via the  $LU$ -factorization of  $A - \sigma B$ , are inaccurate or not computable within reasonable time. The Jacobi-Davidson QZ method [7] has the advantage that it computes with the matrices  $A$  and  $B$  directly and that in principle no inverses or exact solves are needed; furthermore, it poses no restrictions on the matrices  $A$  and  $B$ , so it is also applicable if  $B$  is not symmetric positive semidefinite or if both  $A$  and  $B$  are singular. In section 4, it is shown that the Jacobi-Davidson method with harmonic Petrov values has some favorable properties with respect to purification. If, additionally, a preconditioner is available in the form of an  $LU$ -factorization, the correction equation can be solved efficiently and purification is obtained automatically.

Throughout this paper, it will be assumed that the leftmost finite eigenvalues are wanted. This is a natural assumption in practical situations where the stability of steady states for a number of different parameter values is to be determined, see for instance [1, 9]: not only the leftmost eigenvalue is of interest, but also the eigenvalue(s) close to the leftmost that may become the leftmost for different parameter values. The theory extends readily to problems where the rightmost finite eigenvalues are wanted.

The outline of the paper is as follows. Some properties of generalized eigenvalue problems are described in section 2. In section 3, the Arnoldi method with purification is explained and the new scheme is presented, illustrated by numerical examples. The approach based on the JDQZ method is described in section 4. Section 5 concludes.

## 2 Some properties of generalized eigenvalue problems

Central point of the discussion is the generalized eigenproblem

$$Ax = \lambda Bx, \quad x \neq 0,$$

with  $A, B \in \mathbb{R}^{n \times n}$ ,  $x \in \mathbb{C}^n$  and  $\lambda \in \mathbb{C}$ . Only regular matrix pencils will be considered, i.e. pencils  $(A, B)$  for which  $A - \gamma B$  is singular only for a finite number of  $\gamma \in \mathbb{C}$ . Note that  $B$  is allowed to be singular. The corresponding ordinary eigenproblem is

$$Sx = \tilde{\lambda}x, \quad x \neq 0,$$

with  $S = (A - \sigma B)^{-1}B$  for a  $\sigma$  such that  $A - \sigma B$  is non singular. A generalized eigenpair  $(\lambda, x)$  corresponds to an ordinary eigenpair  $(\tilde{\lambda} = (\lambda - \sigma)^{-1}, x)$  of (1.2). The generalized eigenvalues can be computed via the relation  $\lambda = \tilde{\lambda}^{-1} + \sigma$ .

The eigenspace corresponding to the infinite eigenvalues is the null space  $\mathcal{N}(S)$  of  $S$ :

$$V_\infty = \mathcal{N}(S) = \mathcal{N}(B) = \{x \in \mathbb{R}^n \mid Bx = 0\}.$$

The eigenvectors corresponding to the finite eigenvalues span a real invariant subspace of  $S$  and form a subspace of the range of  $S^{j_s}$ ,  $\mathcal{R}(S^{j_s})$ :

$$V_{finite} \subseteq \mathcal{R}(S^{j_s}) = \{x \in \mathbb{R}^n \mid ((A - \sigma B)^{-1}B)^{j_s}y = x, y \in \mathbb{R}^n\}, \quad (2.1)$$

where  $j_s$  is the size of the largest Jordan block corresponding to the zero eigenvalue of  $S$ . The generalized null space  $\mathcal{G}(S)$  of  $S$  is defined as the complement in  $\mathcal{N}(S^{j_s})$  of  $\mathcal{N}(S)$ :

$$\mathcal{G}(S) = \mathcal{N}(S^{j_s}) \setminus \mathcal{N}(S),$$

It follows that  $\mathbb{R}^n = \mathcal{R}(S^{j_s}) + \mathcal{G}(S) + \mathcal{N}(S)$ . Note that (2.1) becomes an equality if for all finite eigenvalues the algebraic multiplicity is equal to the geometric multiplicity.

It is important to keep in mind that eigenvectors  $v$  corresponding to finite eigenvalues do not necessarily satisfy  $v \perp V_\infty$ . In other words, in general it does not hold that  $\mathcal{R}(S) \perp \mathcal{N}(S)$ . So restricting the search space to  $\mathcal{R}(B)$ , to avoid approximations to infinite eigenvalues, is not effective. Only if  $A$  is block upper triangular and  $B$  is block diagonal it is effective, but then

the problem can also easily be reduced to a smaller problem by considering the non-zero diagonal blocks.

This paper is concerned with block structured generalized eigenvalue problems of the form

$$\begin{bmatrix} K & C \\ C^T & 0 \end{bmatrix} \begin{bmatrix} u \\ p \end{bmatrix} = \lambda \begin{bmatrix} M & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} u \\ p \end{bmatrix}, \quad (2.2)$$

with  $n = m + k$ ,  $C \in \mathbb{R}^{m \times k}$  of full rank, stiffness matrix  $K \in \mathbb{R}^{m \times m}$ , mass matrix  $M = M^T \in \mathbb{R}^{m \times m}$ , velocity  $u \in \mathbb{C}^m$  and pressure  $p \in \mathbb{C}^k$ , that arise in the linearized stability analysis of steady state solutions of the Navier-Stokes equations [1]. The corresponding ordinary eigenproblem is

$$\begin{bmatrix} S_1 & 0 \\ S_2 & 0 \end{bmatrix} \begin{bmatrix} u \\ p \end{bmatrix} = \tilde{\lambda} \begin{bmatrix} u \\ p \end{bmatrix}, \quad S_1 \in \mathbb{R}^{m \times m}, S_2 \in \mathbb{R}^{k \times m},$$

and, as is also noted in [14], this leads to the reduced problem

$$S_1 u = \tilde{\lambda} u, \quad S_1 \in \mathbb{R}^{m \times m}. \quad (2.3)$$

If  $(\tilde{\lambda}, u)$  is an exact eigenpair of  $S_1$ , then for nonzero  $\tilde{\lambda}$ ,  $(\tilde{\lambda}, [u^*, p^*]^*)$  with  $p = \tilde{\lambda}^{-1} S_2 u$  is an exact eigenpair of  $S$ . It can be shown [14, section 2.3] that  $\dim(\mathcal{N}(S_1)) = k$ ,  $\dim(\mathcal{R}(S_1)) = m - k$  and

$$u \in \mathcal{N}(S_1) \quad \Leftrightarrow \quad \begin{bmatrix} u \\ 0 \end{bmatrix} \in \mathcal{G}.$$

Hence, by reducing the problem to (2.3), the geometric multiplicity of the  $k$  eigenvalues  $\tilde{\lambda} = 0$  is reduced from 2 to 1.

### 3 Arnoldi methods with purification

In section 3.1, the implicitly restarted  $B$ -orthogonal Arnoldi method will be described. In section 3.2, it will be shown how this method can be improved by exploiting the specific structure of the generalized eigenproblem. A new strategy, based on this known but previously not used fact, for the computation of a few leftmost eigenvalues will be presented in section 3.3, followed by numerical examples in section 3.4.

#### 3.1 Implicitly restarted $B$ -orthogonal Arnoldi methods

##### 3.1.1 $B$ -orthogonal Arnoldi

The  $B$ -orthogonal Arnoldi method is the standard Arnoldi method with the usual inner product  $(x, y) = x^T y$  replaced by the semi-inner product  $(x, y)_B = x^T B y$ . The  $B$ -orthogonal Arnoldi method constructs a  $B$ -orthonormal basis  $v_1, \dots, v_{k+1}$  for the Krylov subspace

$$\mathcal{K}^{k+1}(S, v_1) = \text{span}(v_1, S v_1, \dots, S^k v_1),$$

where  $S = (A - \sigma B)^{-1} B$ . The basis vectors are related by

$$S V_k = V_k H_k + h_{k+1,k} v_{k+1} e_k^T = V_{k+1} \underline{H}_k, \quad V_{k+1}^T B V_{k+1} = I, \quad (3.1)$$

where  $V_k = [v_1, \dots, v_k] \in \mathbb{R}^{n \times k}$  and  $H_k \in \mathbb{R}^{k \times k}$  and  $\underline{H}_k = [H_k^T, h_{k+1,k} e_k^T]^T \in \mathbb{R}^{(k+1) \times k}$  are upper Hessenberg<sup>1</sup>. Relation (3.1) characterizes a  $k$ -step Arnoldi factorization. Like in the standard Arnoldi method, approximate eigenpairs  $(\theta_i, V_k y_i)$ , i.e. Ritz pairs, can be computed from eigenpairs  $(\theta_i, y_i)$  of  $H_k$ .

<sup>1</sup>Barred identifiers  $\underline{H}_k$  are elements of  $\mathbb{R}^{(k+1) \times k}$ , whereas  $H_k \in \mathbb{R}^{k \times k}$ .

The usual criterion for convergence of a Ritz pair  $(\theta, V_k x)$  with  $H_k x = \theta x$  is derived from the relation

$$SV_k x = V_k H_k x + h_{k+1,k} v_{k+1} e_k^T x = \theta V_k x + h_{k+1,k} v_{k+1} e_k^T x.$$

If  $||h_{k+1,k} v_{k+1} e_k^T x||$  is smaller than a given tolerance  $\tau$ , the Ritz pair  $(\theta, V_k x)$  is said to be converged. It follows that if the  $v_i$  are orthonormalized in the 2-norm, it suffices to inspect  $|h_{k+1,k} e_k^T x|$ . Since for  $B$ -orthogonal Arnoldi the  $B$ -inner product is used, the convergence criterion becomes

$$|h_{k+1,k} e_k^T x| \cdot ||v_{k+1}||_2 < \tau. \quad (3.2)$$

In [14] and [16], the use of the semi-inner product is motivated by the fact that the  $B$  inner product is not affected by components of  $v_i$  in the null space of  $B$ ,  $\mathcal{N}(B)$ , and hence  $H_k$  is independent of components in  $\mathcal{N}(S)$ . Note, however, that  $H_k$  can be corrupted by components in the generalized null space  $\mathcal{G}(S)$ . Moreover, *because* the  $B$ -inner product is not affected by components in the null space of  $B$ , there is no reason to assume that components of  $v_i$  in the null space of  $B$  will not grow; for  $z \in \mathcal{N}(B)$ ,  $x \in \mathbb{R}^n$  and  $\alpha \in \mathbb{R}$ , one has  $||x||_B = ||x + \alpha z||_B$ . As a consequence, the Ritz vector  $V_k y_i$  will be spoiled with error components in  $\mathcal{N} + \mathcal{G}$  (see also [14, Sect. 4.1] and [16, Sect. 2.3]).

The presence of components in  $\mathcal{N} + \mathcal{G}$  in the Arnoldi basis may not only cause spurious eigenvalues and inaccurate Ritz vectors, it may also hamper convergence to the wanted eigenvalues. Purification techniques aim at eliminating the components in  $\mathcal{N} + \mathcal{G}$  from the Arnoldi vectors, with the following three goals:

- removal of spurious eigenpair approximations;
- improvement of wanted eigenpair approximations by removing  $\mathcal{N} + \mathcal{G}$  components from the Ritz vectors;
- increase of the speed of convergence.

Following [14], the notion of purification can be used in several ways, but the idea boils down to eliminating components in  $\mathcal{N} + \mathcal{G}$  of a vector  $x$  by applying  $S^{j_s}$  to it, either explicitly or implicitly. In exact arithmetic, the effect is that  $S^{j_s} x \in \mathcal{R}(S^{j_s})$ , i.e.  $S^{j_s} x$  is in the wanted eigenspace. See [6] and [16] for the first occurrences of the term purification.

### 3.1.2 Implicitly restarted $B$ -orthogonal Arnoldi with purification

In [14, Sect. 3.2], an implicitly restarted Arnoldi method with  $B$ -inner product is proposed. The method, see Alg. 1, reduces the corruption of  $H_k$  by components in  $\mathcal{N}$  and  $\mathcal{G}$  significantly (after the implicit restart), and only requires one additional purification step of the Ritz vectors. The result

---

#### Algorithm 1 Implicitly restarted $B$ -orthogonal Arnoldi with purification

---

- 1: Choose an initial vector  $v_1 \leftarrow S^2 v_1$
  - 2: Do  $k + 1$  steps of  $B$ -orthogonal Arnoldi to compute  $V_{k+2}$  and  $\underline{H}_{k+1}$
  - 3: Compute the  $QR$ -factorization  $\underline{H}_{k+1} = \underline{Q}_{k+1} R_{k+1}$
  - 4: Implicitly restart:  $W_{k+1} = V_{k+2} \underline{Q}_{k+1}$ ,  $\underline{G}_k = R_{k+1} \underline{Q}_k$
  - 5: Compute eigenpairs  $(\theta_i, y_i)$  of the upper  $k \times k$  part of  $\underline{G}_k$
  - 6: Purify the Ritz vectors:  $x_i = S(W_k y_i) = W_{k+1} \underline{G}_k y_i$
  - 7: The eigen approximations for the generalized problem are  $(1/\theta_i + \sigma, x_i)$
- 

of the implicit restart in step 4, is that the Arnoldi vectors in  $W_{k+1}$  and the upper Hessenberg  $\underline{G}_k$  are the same as the ones that would have been computed with starting vector  $Sv_1/||Sv_1||_B$ . In other words, the implicit restart removes the  $\mathcal{N}$  part from  $V_{k+2}$  and the  $\mathcal{G}$  part from  $\underline{H}_{k+1}$ , and it maps the  $\mathcal{G}$  part from  $V_{k+2}$  to the  $\mathcal{N}$  part of  $W_{k+1}$ . Note that because of the  $B$ -inner product,  $\underline{H}_{k+1}$  and  $\underline{G}_k$  are free of contributions of components in  $\mathcal{N}$ . The second purification, in step 6, removes the  $\mathcal{N}$  part from the Ritz vector (the  $\mathcal{G}$  part was already removed by the implicit

restart). The method can still fail due to corruption of  $\underline{H}_{k+1}$  by rounding errors, but this can be detected by inspecting  $\|R_k^{-1}\|_2$  [14, Thm. 4]: if  $\|R_k^{-1}\|_2$  is large and growing for successive values of  $k$ , spurious Ritz values may be computed. Secondly, purification of the Ritz vector  $W_k y_i$  may fail if the corresponding Ritz value  $\theta_i$  is small, i.e.  $\theta_i \sim \epsilon \|S\|$ .

### 3.2 Exploiting the structure of $Ax = \lambda Bx$

In [14, p. 670], [10, p. 8] and [2, p. 1313] it is concluded that the reduced problem

$$S_1 u = \tilde{\lambda} u, \quad S_1 \in \mathbb{R}^{m \times m},$$

see also (2.3), is only of theoretical interest, because  $S_1$  and  $S_2$  depend on blocks in  $A^{-1}$  which are unlikely to be known. However, matrix vector multiplications with  $S_1$ , the only operation with  $S_1$  that is required by the Arnoldi algorithm, and with  $S_2$ , can easily be performed by making use of the available multiplication with  $S$ . Note that in practical situations also  $S$  is not available explicitly and that matrix vector multiplications with  $S$  are for instance implemented using the  $LU$ -factorization of  $A$ .

**Theorem 3.1.** *Let  $S \in \mathbb{R}^{n \times n}$  have the block structure*

$$\begin{bmatrix} S_1 & 0 \\ S_2 & 0 \end{bmatrix},$$

with  $S_1 \in \mathbb{R}^{m \times m}$ ,  $S_2 \in \mathbb{R}^{k \times m}$ , and let  $P = [I_m, 0]^T \in \mathbb{R}^{n \times m}$ ,  $Q = [0, I_k]^T \in \mathbb{R}^{n \times k}$  with  $I_m \in \mathbb{R}^{m \times m}$  an identity matrix of dimension  $m$ . Then for  $x \in \mathbb{C}^m$ ,

$$\begin{aligned} S_1 x &= P^T S P x, \\ S_2 x &= Q^T S P x. \end{aligned}$$

*Proof.* The results follow immediately from the identities  $S_1 = P^T S P$  and  $S_2 = Q^T S P$ .  $\square$

The operations with  $P$  and  $Q$  in Theorem 3.1 can be performed very efficiently and hence with virtually no additional costs the Arnoldi method can be applied to  $S_1$ . This leads to Alg. 2, a modification of Alg. 1. The Arnoldi basis vectors have length  $m < n$ , which reduces the costs of orthogonalization (although usually the costs of operations with  $S$  are dominant). In step 1, only a single explicit purification of the initial vector is needed. Furthermore, the  $B$ -inner product, that was used for its purifying property, and the purification in step 6, are no longer needed, because the implicit restart removes all corruption by components in  $\mathcal{N}$  from  $V_{k+2}$  and  $H_{k+1}$ . On the other hand, to recover the eigenvectors of the original problem, an additional multiplication with  $S_2$  is needed.

---

#### Algorithm 2 Implicitly restarted Arnoldi for $S_1$

---

- 1: Choose an initial vector  $v_1 \leftarrow S_1 v_1 \in \mathbb{R}^m$
  - 2: Do  $k + 1$  steps of Arnoldi with  $S_1$  to compute  $V_{k+2}$  and  $\underline{H}_{k+1}$
  - 3: Compute the  $QR$ -factorization  $\underline{H}_{k+1} = \underline{Q}_{k+1} R_{k+1}$
  - 4: Implicitly restart:  $W_{k+1} = V_{k+2} \underline{Q}_{k+1}$ ,  $\underline{G}_k = R_{k+1} \underline{Q}_k$
  - 5: Compute eigenpairs  $(\theta_i, y_i)$  of the upper  $k \times k$  part of  $\underline{G}_k$
  - 6: Compute  $p_i = \theta_i^{-1} S_2 x_i$
  - 7: The eigen approximations for the generalized problem are  $(1/\theta_i + \sigma, [x_i^*, p_i^*]^*)$
- 

#### 3.2.1 Improved rounding error analysis

The most important consequence of Theorem 3.1, however, is that the results of the error analysis in [14, section 5] improve considerably. Following the notation and assumptions there, let  $P_{\mathcal{R}_1}$  and

$P_{\mathcal{N}_1}$  be normalized projectors that map a vector into  $\mathcal{R}_1 = \mathcal{R}(S_1)$  and  $\mathcal{N}_1 = \mathcal{N}(S_1)$  respectively, so  $x \in \mathbb{C}^m$  can be decomposed uniquely as  $x = P_{\mathcal{R}_1}x + P_{\mathcal{N}_1}x$ . Note that  $P_{\mathcal{N}_1}S_1 = 0$ . The computed Arnoldi vectors satisfy

$$\begin{aligned} h_{j+1,j}v_{j+1} &= Sv_j - \sum_{i=1}^j h_{ij}v_i + \psi_j, \\ h_{ij} &= v_i^T Sv_j + \delta_{ij}, \\ v_i^T v_j &= \begin{cases} 1 + \gamma_{ij}, & j = i \\ \gamma_{ij} & j = 1, \dots, i+1, j \neq i. \end{cases} \end{aligned}$$

In block form, the round-off errors  $\|\Psi_{k+1}\|_2$ ,  $\|\Gamma_{k+1}\|_2$  and  $\|\Delta_k\|_2$  for the  $k$ -step Arnoldi factorization are given by the following relations:

$$V_{k+1}\underline{H}_k = SV_k + \Psi_{k+1}, \quad (3.3)$$

$$V_{k+1}^T V_{k+1} = I + \Gamma_{k+1}, \quad (3.4)$$

$$\underline{H}_k = V_{k+1}^T SV_k + \Delta_k. \quad (3.5)$$

**Result 3.2.** *The  $\mathcal{N}_1$  component in  $v_j$  may increase as  $j$  increases.*

*Proof.* Repeating the proof in [14, section 4.1] leads to

$$\begin{aligned} h_{j,j+1}P_{\mathcal{N}_1}v_{j+1} &= P_{\mathcal{N}_1}S_1v_j - \sum_{i=1}^j h_{ij}P_{\mathcal{N}_1}v_i + P_{\mathcal{N}_1}\psi_j \\ &= -\sum_{i=1}^j h_{ij}P_{\mathcal{N}_1}v_i + P_{\mathcal{N}_1}\psi_j. \end{aligned}$$

There is no reason to assume that  $\|P_{\mathcal{N}_1}v_{j+1}\|$  does not increase.  $\square$

The improvement over the result in [14, section 4.1] is that for Arnoldi applied to  $S$ , there may be an increase of both components in  $\mathcal{N}$  and  $\mathcal{G}$ , while here there only may be a smaller increase of components in  $\mathcal{N}_1$ .

The following result shows the improved effect of the implicit purification via  $x_j = S(V_k z_j) = V_{k+1}\underline{H}_k z_j$ , where  $z_j$  is an eigenvector of  $\underline{H}_k z_j$ . Although this purification is not needed in Alg. 2, as will become clear in result 3.4 and theorem 3.5, it is included here, however, to show that the relative contributions of the  $\mathcal{N}_1$  components are smaller than in the results in [14, section 4.2].

**Result 3.3.** *The purification operation  $x_j = V_{k+1}\underline{H}_k z_j$  produces an approximate eigenvector with no  $\mathcal{N}_1$  component. This step may fail if  $|\theta_j^{-1}| \gg \epsilon_M^{-1}$ , where  $\epsilon_M$  is the machine precision number.*

*Proof.* From the proof in [14, section 4.2], it follows that the purified  $x_j$  computed by  $x_j = V_{k+1}\underline{H}_k z_j$  with  $\|z_j\|_2 = 1$  satisfies

$$\begin{aligned} P_{\mathcal{N}_1}x_j &= P_{\mathcal{N}_1}S_1V_k z_j + P_{\mathcal{N}_1}\xi_j \\ &= P_{\mathcal{N}_1}\xi_j, \end{aligned} \quad (3.6)$$

with

$$\|\xi_j\|_2 \leq 3k^{3/2}\|V_{k+1}\|_F\|A^{-1}\|_2\epsilon_M + \|\Psi_{k+1}\|_2 + O(\epsilon_M^2).$$

If  $\|z_j\|_2 = 1$  (note that  $H_k z_j = \theta_j z_j$ ), then  $\|V_k z_j\|_2 \simeq 1$  and  $\|x_j\|_2 \simeq \theta_j$  and hence relative contributions of the  $\mathcal{N}_1$  components in  $x_j$  are obtained by dividing (3.6) by  $\theta_j$ . If  $\theta_j$  is small, these relative contributions become large and purification may fail. If  $|\theta_j^{-1}|\|\xi_j\|_2 \ll 1$ , then the  $\mathcal{N}_1$  component in  $x_j$  is removed.  $\square$

**Result 3.4.** *One implicit restart of Arnoldi produces a  $\underline{G}_k$  that is not corrupted by  $\mathcal{N}_1$  components, and a  $W_{k+1}$  that has no  $\mathcal{N}_1$  component. This step may fail if  $\|R_{k+1}^{-1}\|_2 \gg \epsilon_M^{-1}$ .*

*Proof.* Repeating the proof in [14, section 4.4] leads to

$$\|P_{\mathcal{N}_1}W_{k+1}\|_2 \leq \|P_{\mathcal{N}_1}\Xi_{k+1}\|_2 + O(\epsilon_M^2),$$

with

$$\begin{aligned} \|\Xi_{k+1}\|_2 &\leq (k+2)^{3/2}\|V_{k+2}\|_F\epsilon_M \\ &\quad + (\omega\|V_{k+1}\|_2\|A^{-1}\|_2\epsilon_M + \|\Psi_{k+2}\|_2)\|R_{k+1}^{-1}\|_2 + O(\epsilon_M^2), \end{aligned}$$

and  $\omega = O(1)$ . If  $\|R_{k+1}^{-1}\|_2$  is small,  $W_{k+1}$  has no significant components in  $\mathcal{N}_1$ . If  $\|R_{k+1}^{-1}\|_2$  is large, then  $\|\Xi_k\|_2 \gg \epsilon_M$  and  $W_{k+1}$  can have components in  $\mathcal{N}_1$ . Consequently,  $G_k$  is corrupted by the components in  $\mathcal{N}_1$  and may cause spurious Ritz values.  $\square$

Compared to the results in [14, section 4.4], the corruption in  $W_{k+1}$  and  $G_k$  is decreased.

The following theorem shows that, as a Ritz pair  $(\tilde{\lambda}, x)$  with not too small  $\tilde{\lambda}$  converges to an eigenpair of  $S_1$ , it is purified automatically. This is consistent with results 3.3 and 3.4, and also explains why implicit purification of converged Ritz pairs (step 6 in Alg. 1) is not needed.

**Theorem 3.5.** *Let  $(\tilde{\lambda}, x)$  be a converged Ritz pair of  $S_1$ , with  $r = S_1x - \tilde{\lambda}x$  and  $\|r\|_2 < \epsilon$ . Then  $\|P_{\mathcal{N}_1}x\|_2 \leq \epsilon/\tilde{\lambda}$ .*

*Proof.* Write  $\tilde{\lambda}P_{\mathcal{N}_1}x = P_{\mathcal{N}_1}S_1x - P_{\mathcal{N}_1}r$  and note that  $P_{\mathcal{N}_1}S_1x = 0$ .  $\square$

Although failure of IRA if  $\|R_k^{-1}\|_2$  is large is still possible, the results above show that the (growth of the) corruption by  $\mathcal{N}_1$  components is reduced. The rounding errors made during the orthogonalization phase are also reduced, because the standard inner product is used instead of the  $B$ -inner product, and hence no additional multiplications with  $B$  are needed.

### 3.2.2 Numerical example

To illustrate the new results, the growth of  $\|\Psi_{k+1}\|_2$  (see (3.3)) for  $S$  and  $S_1$  is compared. Figure 1 shows  $\|\Psi_{k+1}\|_2$  at every Arnoldi iteration for the example matrix pencil taken from [14, Sect. 3.3]. For  $S$ , the  $B$ -orthogonal Arnoldi method is used, while for  $S_1$  Arnoldi with the usual inner product is used. For both cases, the initial vector  $v_1$ , with all entries equal to one, was purified using  $v_1 \leftarrow S^2v_1$  and  $v_1 \leftarrow S_1v_1$  respectively. It is clear that the growth of  $\|\Psi_{k+1}\|_2$  is much smaller for  $S_1$ . The growth of  $\|\Psi_{k+1}\|_2$  for  $S$  can be explained as follows: let  $w_k$  be the new Arnoldi vector in iteration  $k$ , just after orthogonalization against  $V_k$ , but before normalization. The  $B$ -inner product neglects any components in  $\mathcal{N}(B)$ , but these components are normalized with the same factor  $h_{k+1,k} = \|w_k\|_B$ . If  $h_{k+1,k} < 1$ , then these components increase in 2-norm. This may lead to an increase of  $\|\Psi_{k+1}\|_2$ . Typical values of  $h_{k+1,k}$  in this example were of order  $O(10^{-4})$ . An explanation for the apparent stagnation of the growth of  $\|\Psi_{k+1}\|_2$  at some iterations may be: the new Arnoldi vector is computed as  $Sv_k$  and  $S_1v_k$  respectively, which is in fact an explicit purification of  $v_k$ . Combined with a not too small  $h_{k+1,k}$ , this will cause only a limited increase.

A large  $\|\Psi_{k+1}\|_2$  may not only prevent the implicit restart with zero shift from purifying the factorization, it also reduces the effect of the implicit purification via  $x_j = V_{k+1}\underline{H}_k z_j$ , as can be deduced from results 3.3 and 3.4 and their equivalents in [14]. With this in mind, the choice for Arnoldi with  $S_1$  is obvious.

## 3.3 A new strategy

### 3.3.1 Implicitly restarted Arnoldi with deflation

It is not clear from [14] how the idea of the implicit restart with shift  $\sigma_0 = 0$  (Alg. 1) is incorporated with the implicitly restarted Arnoldi method with deflation [11, 22]. The IRA method starts with a  $k$ -step Arnoldi factorization  $SV_k = V_kH_k + h_{k+1,k}v_{k+1}e_k^T$ . Then, until convergence, the following steps are iterated:






Figure 1: The size of  $\|\Psi_{k+1}\|_2$  for  $B$ -orthogonal Arnoldi applied to  $S = (A - 60B)^{-1}B$ , and Arnoldi applied to  $S_1$ .

1. Compute the Ritz values  $\theta_i$ , i.e. the eigenvalues of  $H_k$  and split them in a set of wanted Ritz values  $\{\theta_1 \dots \theta_j\}$  and unwanted Ritz values  $\{\sigma_1 \dots \sigma_p\}$ , with  $k = j + p$ .
2. Apply  $p$   $QR$ -steps to  $H_k$  with shifts  $\sigma_i$  to remove the unwanted Ritz values.
3. Extend the  $j$ -step Arnoldi factorization to a  $k$ -step Arnoldi factorization.

Like in Alg. 2, the idea now is to implicitly restart with  $\sigma_0 = 0$  just before the computation of the Ritz values in step (1), i.e. just after the extension of the Arnoldi factorization. Any detected spurious Ritz values can be removed by including these as shifts for the implicit restarts. The algorithm is summarized in Alg. 3. For details about the implementation of implicit shifts, deflation and the locking procedure, the reader is referred to [11, 22, 23].

---

**Algorithm 3** Implicitly restarted Arnoldi for  $S_1$  with purification and deflation

---

- 1: Choose an initial vector  $v_1 \leftarrow S_1 v_1$
  - 2: Do  $k + 1$  steps of Arnoldi to compute  $V_{k+2}$  and  $\underline{H}_{k+1}$
  - 3: **while** not all converged **do**
  - 4: Purify by applying one restart with  $\sigma = 0$ :  $[V_{k+1}, \underline{H}_k] = \text{purify}(V_{k+2}, \underline{H}_{k+1})$
  - 5: Compute  $\lambda(H_k)$  and lock converged wanted Ritz values
  - 6: Select  $p$  shifts  $\sigma_1, \dots, \sigma_p$
  - 7: Apply  $p$  implicit shifts to compute the  $(k - p)$  step Arnoldi factorization  $S_1 V_{k-p} = V_{k-p+1} \underline{H}_{k-p}$
  - 8: Extend  $S_1 V_{k-p} = V_{k-p+1} \underline{H}_{k-p}$  to  $S_1 V_{k+1} = V_{k+2} \underline{H}_{k+1}$
  - 9: **end while**
- 

### 3.3.2 Exploiting transformations to improve selection and convergence

Besides the shift-and-invert transformation  $T_{SI}(A, B, \sigma) = (A - \sigma B)^{-1}$ , the generalized Cayley transformation

$$T_C(A, B, \alpha_1, \alpha_2) = (A - \alpha_1 B)^{-1}(A - \alpha_2 B) = B + (\alpha_1 - \alpha_2)T_{SI}, \quad \alpha_1, \alpha_2 \in \mathbb{R} \quad (3.7)$$

with  $\alpha_1 < \alpha_2$  and  $\alpha_1 \neq \lambda_i, i = 1, \dots, n$ , can be used for problems of the form (2.2), see [1, 2, 10]. The eigenvalues  $\mu_i$  of  $T_C$  are related to the eigenvalues of  $(A, B)$  by the relation  $\mu_i = (\lambda_i - \alpha_1)^{-1}(\lambda_i - \alpha_2)$  and the infinite eigenvalues are transformed to 1. Eigenvalues close to  $\alpha_1$  are mapped to eigenvalues far from the unit circle, while eigenvalues close to  $\alpha_2$  are mapped to eigenvalues with small magnitude. The property that is of most use is that eigenvalues with  $\text{Re}(\lambda_i) < (\alpha_1 + \alpha_2)/2$  are mapped outside the unit circle, while eigenvalues with  $\text{Re}(\lambda_i) > (\alpha_1 + \alpha_2)/2$  are mapped inside the unit circle. The modified Cayley transformation is defined by

$$T_M(A, B, \alpha_1, \alpha_2, \alpha_3) = \begin{bmatrix} K - \alpha_1 M & C \\ C^T & 0 \end{bmatrix}^{-1} \begin{bmatrix} K - \alpha_2 M & \alpha_3 C \\ \alpha_3 C^T & 0 \end{bmatrix}, \quad (3.8)$$

and has the same properties as the generalized Cayley transform, except that the infinite eigenvalues are transformed to  $\alpha_3$  [1, 2, 10].

In [10], an algorithm is described where Cayley-transformations are combined with shift-and-invert Arnoldi. The algorithm is based on the hybrid algorithm presented in [1, section 2.3] and consists of two phases. In the first phase, an  $r$ -step Arnoldi factorization is computed using  $B$ -orthogonal Arnoldi with purification. The corresponding  $r$  Ritz values are used to determine the parameters  $\alpha_1, \alpha_2 \in \mathbb{R}$  of the (modified) Cayley transform  $T_C$ . In the second phase, implicitly restarted  $B$ -orthogonal Arnoldi with purification is applied to  $T_C$  to compute the wanted eigenvalues. The parameters  $\alpha_1, \alpha_2$  are updated during the restarts.

The Ritz values that are computed in phase 1 may not have converged (moreover, in [10] there is *no* convergence testing for the Ritz pairs of phase 1, to avoid accepting wrong eigenvalues), and there may be spurious Ritz values as well. Also in the second phase spurious Ritz values may be computed, that make the determination of  $\alpha_1, \alpha_2$  more difficult. In [10] a selection strategy is used to deal with spurious Ritz values. The approach presented here makes such a strategy unnecessary and also reduces the number of different Cayley-transformations needed.

Assume that the  $k = 2$  leftmost eigenvalues of (2.2) are wanted (complex conjugate pairs counted as one eigenvalue), including any eigenvalues with negative real part. The algorithm is readily adjustable for any number of wanted eigenvalues.

Simply computing the leftmost eigenvalues of  $S = T_{SI}(A, 0) = A^{-1}B$  is not advisable for several reasons. Firstly, even if there are eigenvalues with negative real part, the process will most likely be disturbed by spurious Ritz values, as has been explained in the previous sections. Secondly, the leftmost eigenvalues of  $S$  do not necessarily correspond to the leftmost eigenvalues of  $(A, B)$ . The extremal eigenvalues  $\tilde{\lambda}_i$  of  $S$ , that correspond to the eigenvalues  $\lambda_i = 1/\lambda_i$  of  $(A, B)$ , however, can be computed safely, efficiently and accurately with IRA. These eigenvalues, sorted in increasing real part order, that also not necessarily are the leftmost eigenvalues of  $(A, B)$ , can be used to compute  $\alpha_1, \alpha_2$  for the modified Cayley transform:

- If  $\text{Im}(\lambda_1) = 0$ , then  $\alpha_1 = \lambda_1 + \frac{\text{Re}(\lambda_2) - \lambda_1}{2}$ .
- If  $\text{Im}(\lambda_1) \neq 0$ , then  $\alpha_1 = \lambda_1$ .
- In both cases,  $\alpha_2 = 2 \times \text{Re}(\lambda_2) - \alpha_1$ .

With these choices for  $\alpha_1, \alpha_2$ , eigenvalues with  $\text{Re}(\lambda_i) < \text{Re}(\lambda_2)$  correspond to eigenvalues  $\tilde{\lambda}_i$  of  $S_M = T_M(A, B, \alpha_1, \alpha_2, 0)$  with  $|\tilde{\lambda}_i| > 1$ , while eigenvalues with  $\text{Re}(\lambda_i) > \text{Re}(\lambda_2)$  are transformed inside the unit circle. Hence also any missed eigenvalues between  $\lambda_1$  and  $\lambda_2$  correspond to eigenvalues  $\mu_i$  of  $S_M$  with  $|\mu_i| > 1$ . The eigenvalues of  $S_M$  with largest magnitude can again be computed by IRA and because the infinite eigenvalues are transformed to  $\alpha_3 = 0$ , there is virtually no danger that spurious Ritz values will be selected as wanted eigenvalue. As soon as eigenvalues inside the unit circle are computed, it can be safely concluded that the leftmost eigenvalues (including the eigenvalues with negative real part) are found. The strategy is shown in Alg. 4.

The strategy consists of two phases: in phase 1 (step 1-2), the largest eigenvalues (in magnitude) of  $S$  are computed. Phase 2 (step 3-6) checks for any missed eigenvalues using the Cayley transformation. In step 1, a larger number  $r$  will increase the chance of computing the leftmost

---

**Algorithm 4** Strategy for computing the 2 left most eigenvalues of  $(A, B)$

---

**PHASE 1**

- 1: Compute the  $r \geq 2$  largest eigenvalues  $\tilde{\lambda}_i$  of  $S_1$  with Alg. 3
- 2: Order  $\lambda_i = 1/\tilde{\lambda}_i$ ,  $i = 1, \dots, r$  by increasing real part

**PHASE 2**

- 3: Determine  $\alpha_1$  and  $\alpha_2$ , and  $\alpha_3 = 0$
  - 4:  $S_M = T_M(A, B, \alpha_1, \alpha_2, \alpha_3)$
  - 5: Compute the largest 2 eigenvalues  $\mu_i$  of  $S_{M1}$  with Alg. 3
  - 6: The eigenvalues of  $(A, B)$  are  $\lambda_i = \frac{\alpha_1 \mu_i - \alpha_2}{\mu_i - 1}$
- 

eigenvalue already in this phase. In step 4, one could also take  $\alpha_1 = 0$ , but because any missed eigenvalues are expected to be close to  $\lambda_1$ , this is not preferred. Additional verification of any missed eigenvalues can be done by choosing new  $\alpha_1, \alpha_2$  based on the eigenvalues found in step 7 to compute the largest eigenvalues of the new  $S_M$ , or by using techniques described in [15].

The difference with existing approaches is that in the determination of  $\alpha_1, \alpha_2$  rather accurate eigenvalue approximations are used, with as advantages that fewer updates of  $S_M$  are needed and that the risk of missing eigenvalues is reduced. Furthermore, the possible disturbance by spurious Ritz values is reduced by first computing only the largest eigenvalues of  $S_1$ . Note that with the choice  $\alpha_3 = 0$ ,  $S_M$  can be reduced to  $S_{M1}$  in the same way as  $S$  to  $S_1$ , as described in section 3.2. If  $(\mu, u)$  is an exact eigenpair of  $S_{M1}$ , then for nonzero  $\mu$ ,  $(\mu, [u^*, p^*]^*)$  with  $p = \mu^{-1} S_{M2} u$  is an exact eigenpair of  $S_M$ . If  $(\mu, [u^*, p^*]^*)$  is an eigenpair of the modified eigenvalue problem (3.8), then  $(\frac{\alpha_1 \mu_i - \alpha_2}{\mu_i - 1}, [u^*, q^*]^*)$ , with  $q = (\mu - \alpha_3)/(\mu - 1)p$ , is an eigenpair of the original generalized eigenvalue problem (2.2), provided  $\mu \notin \{1, \alpha_3\}$  (see [8, 10]).

It may seem that there is no advantage in using  $S_M = T_M(\alpha_1, \alpha_2, 0)$  instead of  $S$ , since in exact arithmetic, due to shift-invariance of Krylov subspaces, Arnoldi for  $S$  and  $S_M$  produces the same eigenvalue estimates of  $(A, B)$  [15, lemma 2.5]. However, when using Arnoldi for  $S$ , the spurious Ritz values may be hard to distinguish from wanted leftmost Ritz values, as both may be close to zero, while when using Arnoldi for  $S_M$ , the spurious Ritz values (near zero) are clearly separated from the wanted Ritz values (magnitude larger than 1).

### 3.4 Realistic examples

The strategy in Alg. 4 is applied to two large scale examples. The first example is the stability analysis of the flow over a backward facing step, a well known benchmark problem from fluid dynamics [9]. The second examples is the flow in a driven cavity [4, section 7.1.3]. When referring to (finite) eigenvalues  $\lambda_i$ , it is assumed that the  $\lambda_i$  are sorted in increasing real part order, i.e.  $\lambda_1$  is the leftmost eigenvalue. For more information about the bifurcation analysis of such nonlinear systems, see [3].

The method `eigs` of Matlab 6.5, which is a wrapper around ARPACK [12], is used in all experiments. The stopping criterion is  $\tau = 10^{-6}$  and the size of the Arnoldi factorization is  $k = 20$ .

#### 3.4.1 Flow over a backward facing step

The matrices  $A$  and  $B$  with  $n = m + p = 21,730 + 7,872 = 29,602$ , were obtained using the package IFISS [18]. Table 1 shows statistics for Alg. 4 with  $r = 2$ , both for  $S$  and the reduced problem  $S_1$ . The leftmost eigenvalues  $\lambda_1 = 6.04 \cdot 10^{-2}$  and  $\lambda_{2,3} = 7.97 \cdot 10^{-2} \pm i1.92 \cdot 10^{-2}$  were already found in the first phase of the algorithm: the validation in phase 2 did not result in new eigenvalues. Although the running times for both the reduced and the unreduced problem are equal, as expected, the residuals are better for the reduced problem. The claim in [9], that the steady state flow at a Reynolds number  $Re = 800$  is stable, is confirmed by the results.

Table 1: Statistics for Alg. 4 for the flow over a backward facing step (section 3.4.1): number of restarts, time, found eigenvalues and residuals after each phase.

	reduced		unreduced	
	phase 1	phase 2	phase 1	phase 2
#restarts	3	2	3	3
time (s)	118	95	120	97
eigenvalues	$\lambda_1, \lambda_{2,3}$	$\lambda_1, \lambda_{2,3}$	$\lambda_1, \lambda_{2,3}$	$\lambda_1, \lambda_{2,3}$
$\max_i \ Ax_i - \lambda_i Bx_i\ $	$1 \cdot 10^{-12}$	$1 \cdot 10^{-12}$	$9 \cdot 10^{-11}$	$9 \cdot 10^{-11}$

Table 2: Statistics for Alg. 4 for the driven cavity (section 3.4.2): number of restarts, time, found eigenvalues and residuals after each phase.

	$r = 2$		$r = 5$	
	phase 1	phase 2	phase 1	phase 2
#restarts	1	4	6	4
time (s)	33	112	106	112
eigenvalues	$\lambda_1, \lambda_4$	$\lambda_1, \lambda_{2,3}$	$\lambda_1, \lambda_{4-7}$	$\lambda_1, \lambda_{2,3}$
$\max_i \ Ax_i - \lambda_i Bx_i\ $	$1 \cdot 10^{-16}$	$1 \cdot 10^{-14}$	$1 \cdot 10^{-10}$	$1 \cdot 10^{-14}$

### 3.4.2 Flow in a driven cavity

The matrices  $A$  and  $B$  with  $n = m + p = 8,450 + 1,089 = 9,539$ , were obtained using the package IFISS [18]. Table 2 shows statistics for Alg. 4 with  $r = 2$  and  $r = 5$ , for the reduced problem  $S_1$ . The eigenvalues  $\lambda_1 = 3.21 \cdot 10^{-2}$  and  $\lambda_4 = 1.01 \cdot 10^{-1}$  were found in the first phase of the algorithm. The validation in phase 2 identified the missed eigenvalue pair  $\lambda_{2,3} = 6.20 \cdot 10^{-2} \pm i4.61 \cdot 10^{-1}$ . Increasing  $r$  does not help finding the missed eigenvalue in phase 1, while it increases the running time.

## 4 Jacobi-Davidson methods, preconditioning and purification

If the linear system solves with  $A - \sigma B$  are inaccurate or not computable within reasonable time, the strategy based on the implicitly restarted Arnoldi method is no longer applicable, although an inexact variant could be considered [13]. Here a scheme based on the Jacobi-Davidson QZ method [7] is proposed, that does not require exact solves with  $(A - \sigma B)$ .

The Jacobi-Davidson method [20] combines two principles to compute eigenpairs of eigenvalue problems  $Ax = \lambda x$ . The first principle is to apply a Ritz-Galerkin approach with respect to a subspace spanned by  $v_1, \dots, v_k$ , the search space. The second principle is the computation of a correction orthogonal the current eigenvector approximation. The Jacobi-Davidson method for generalized eigenvalue problems will be briefly explained in sections 4.1 and 4.2. For a more detailed description, the reader is referred to [7, 19, 20].

In section 4.3, it will be shown that when an exact preconditioner is used to solve the correction equation, purification is obtained automatically. In section 4.4, this fact will be combined with other properties of Jacobi-Davidson to obtain an efficient method for the computation of a few selected eigenvalues.

## 4.1 The Jacobi-Davidson method for generalized eigenproblems

Given the generalized eigenvalue problem

$$Ax = \lambda Bx, \quad x \neq 0,$$

with  $A, B \in \mathbb{R}^{n \times n}$ , the Jacobi-Davidson method applies a Petrov-Galerkin condition to compute approximate eigenpairs. If the search space is spanned by  $v_1, \dots, v_k$ , with  $V_k = [v_1, \dots, v_k]$  orthogonal, and the test space is spanned by  $w_1, \dots, w_k$ , with  $W_k = [w_1, \dots, w_k]$  orthogonal, the Petrov-Galerkin condition becomes

$$AV_k s - \theta BV_k s \perp \{w_1, \dots, w_k\}.$$

This leads to the reduced  $k \times k$  system

$$W_k^* AV_k s = \theta W_k^* BV_k s,$$

which can be solved using full space methods like  $QZ$  to compute eigenpair approximations  $(\theta_i, q_i = V_k s_i)$  of (4.1).

Given such an eigenpair approximation  $(\theta_i, q_i)$ , the question is how to expand the search and test space to improve the approximation. With the corresponding residual vector given by

$$r_i = (Aq_i - \theta_i Bq_i),$$

the Jacobi-Davidson method computes a correction  $t \perp q_i$  from the Jacobi-Davidson correction equation

$$(I - z_i z_i^*)(A - \theta_i B)(I - q_i q_i^*)t = -r_i, \quad (4.1)$$

where the test vector  $z_i = \mu Aq_i + \nu Bq_i$  for a suitable pair  $\mu, \nu \in \mathbb{C}$ . The search space is expanded with  $t$  and the test space is expanded with  $\mu At + \nu Bt$ . A Ritz pair is accepted if  $\|r_i\|_2 = \|(Aq_i - \theta_i Bq_i)\|_2$  is smaller than a given tolerance.

## 4.2 Jacobi-Davidson $QZ$

In [7, 19], the Jacobi-Davidson method is extended with deflation. The Jacobi-Davidson  $QZ$  ( $JDQZ$ ) method computes a partial generalized Schur form of the pencil  $(A, B)$ . Let the current approximate partial generalized Schur form be given by

$$AQ_k = Z_k S_k, \quad BQ_k = Z_k T_k,$$

with  $Q_k, Z_k$   $n \times k$  matrices and  $S_k, T_k$  upper triangular  $k \times k$  matrices. The problem of finding the next Schur triple  $(q_{k+1}, z_{k+1}, (\alpha_{k+1}, \beta_{k+1}))$  with  $\theta_{k+1} = \alpha_{k+1}/\beta_{k+1}$  can be rewritten as a deflated generalized eigenvalue problem

$$Q_k^* q_{k+1} = 0, \quad (I - Z_k Z_k^*)(\beta_{k+1} A - \alpha_{k+1} B)(I - Q_k Q_k^*) q_{k+1} = 0,$$

which can be solved by the Jacobi-Davidson method. With the search space represented by the orthogonal matrix  $V$  and the test space by the orthogonal matrix  $W$ , so that  $V^* Q_k = W^* Z_k = 0$ , the reduced system matrices become

$$\begin{aligned} M_A &\equiv W^*(I - Z_k Z_k^*)A(I - Q_k Q_k^*) = W^*AV, \\ M_B &\equiv W^*(I - Z_k Z_k^*)B(I - Q_k Q_k^*) = W^*BV. \end{aligned}$$

The generalized Schur decomposition of  $(M_A, M_B)$  is computed using  $QZ$ :

$$Z_M^* M_A Q_M = S_A, \quad Z_M^* M_B Q_M = S_B.$$

The generalized Schur form is ordered with respect to the target  $\tau$ , and an approximate Petrov triple for (4.2) is obtained as

$$(\tilde{q}, \tilde{z}, (\tilde{\alpha}, \tilde{\beta})) = (VQ_M e_1, WZ_M e_1, (s_{A,11}, s_{B,11})).$$

Given a Petrov triple  $(\tilde{q}, \tilde{z}, (\tilde{\alpha}, \tilde{\beta}))$  for the deflated problem, the corresponding generalized deflated correction equation becomes

$$(I - \tilde{z}\tilde{z}^*)(I - Z_k Z_k^*)(\tilde{\beta}A - \tilde{\alpha}B)(I - Q_k Q_k^*)(I - \tilde{q}\tilde{q}^*)t = -\tilde{r}_i, \quad (4.2)$$

where the residual  $\tilde{r}$  is

$$\tilde{r} = (I - Z_k Z_k^*)(\tilde{\beta}A - \tilde{\alpha}B)(I - Q_k Q_k^*)\tilde{q},$$

and  $Q_k^* t = Z_k^* \tilde{z} = Q_k^* \tilde{q} = 0$ ,  $\tilde{q}^* t = 0$ ,  $\|t\|_2 = 1$ . The search space is expanded with the orthogonal complement of  $t$ , and the test space is orthogonally expanded with  $(I - Z_k Z_k^*)(\mu A + \nu B)(I - Q_k Q_k^*)t$ .

If the correction equation is solved exactly, the Jacobi-Davidson method converges asymptotically quadratically. In fact, the method can be shown to be a Newton scheme. Solving the correction equation exactly may be too expensive in practice and therefore Krylov subspace methods with preconditioning are used to solve the correction equation approximately. With a preconditioner  $K \approx A - \tau B$ , the correction equation (4.2) can be preconditioned by

$$(I - \tilde{z}\tilde{z}^*)(I - Z_k Z_k^*)K(I - Q_k Q_k^*)(I - \tilde{q}\tilde{q}^*).$$

With  $Q_k := [Q_k, \tilde{q}]$ ,  $Z_k := [Z_k, \tilde{z}]$ ,  $Y_k = K^{-1}Z_k$  and  $H_k = Q_k^* Z_k$ , the left preconditioned correction equation becomes

$$(I - Y_k H_k^{-1} Q_k^*)K^{-1}(\beta A - \alpha B)(I - Y_k H_k^{-1} Q_k^*)t = -r, \quad (4.3)$$

where  $r = (I - Y_k H_k^{-1} Q_k^*)K^{-1}\tilde{r}$ .

### 4.3 Purification

Jacobi-Davidson style methods select a new Petrov pair according to some criterion, for instance the leftmost Petrov pair, at every iteration. In the absence of infinite eigenvalues, selecting the leftmost Petrov pair will usually result in convergence to the leftmost eigenvalue, assuming that the initial search space contains components in that direction. In the presence of infinite eigenvalues however, this will no longer be a smart strategy: Petrov values will go to infinity, without a proper mechanism to identify them as infinite eigenvalue approximations.

If the search space is restricted to  $\mathcal{R}(S^j)$ , approximations to infinite eigenvalues can be avoided. Projection of the search space vectors onto  $\mathcal{R}(S^j)$  is not attractive because an orthogonal basis for  $\mathcal{R}(S^j)$  is not cheaply available. The following lemmas are needed for proving theorem 4.6, which states that if an exact preconditioner<sup>2</sup> is used for the correction equation and if the initial search space  $V_0 \subset \mathcal{R}(S^j)$ , then, in exact arithmetic, no spurious eigenvalues are computed during the Jacobi-Davidson process.

**Lemma 4.1.** *Let  $q = ((A - \sigma B)^{-1}B)^j x \in \mathcal{R}(S^j)$  and  $K = A - \tau_0 B$ . Then  $r = (\beta A - \alpha B)q \in \mathcal{R}(BS^{j-1})$ .*

*Proof.* The result follows from some linear algebra:

$$\begin{aligned} (\beta A - \alpha B)q &= \beta((A - \sigma B) + (\sigma - \alpha/\beta)B)q \\ &= \beta((\sigma - \alpha/\beta)Bq + (A - \sigma B)((A - \sigma B)^{-1}B)^j x) \\ &= \beta B((\sigma - \alpha/\beta)q + ((A - \sigma B)^{-1}B)^{j-1} x) \in \mathcal{R}(BS^{j-1}), \end{aligned}$$

where in the last step  $\mathcal{R}(BS^j) \subseteq \mathcal{R}(BS^{j-1})$  is used.  $\square$

<sup>2</sup>In this paper, an exact preconditioner is a preconditioner  $K = A - \tau_0 B$  for which linear systems of the form  $Kx = y$  can be solved exactly, for instance by using an exact  $LU$ -factorization  $LU = K$ .

**Lemma 4.2.** Let  $y = BS^{j-1}x \in \mathcal{R}(BS^{j-1})$  and  $K = A - \tau_0 B$ . Then  $K^{-1}y \in \mathcal{R}(S^j)$ .

*Proof.* With basic linear algebra, one finds

$$\begin{aligned} K^{-1}y &= (A - \tau_0 B)^{-1}y \\ &= (A - \tau_0 B)^{-1}BS^{j-1}x \\ &= (A - \sigma B)^{-1}(I + (\tau_0 - \sigma)B(A - \tau_0 B)^{-1})BS^{j-1}x \in \mathcal{R}(S^j). \end{aligned}$$

□

**Lemma 4.3.** Let  $r \in \mathcal{R}(S^j)$ ,  $K = A - \tau_0 B$ ,  $AQ_k = Z_k S_A$ ,  $BQ_k = Z_k S_B$ ,  $q \in \mathcal{R}(S^j)$ ,  $z = \nu Aq + \mu Bq$ ,  $Y_k = K^{-1}[Z_k, z]$  and  $H_k = [Q_k, q]^* Z_k$ . Then  $(I - Y_k H_k^{-1}[Q_k, q]^*)r \in \mathcal{R}(S)$ .

*Proof.* First note that  $\mathcal{R}(Z_k) = \mathcal{R}(AQ_k) = \mathcal{R}(BQ_k)$ . It follows from lemma 4.1 that  $z \in \mathcal{R}(BS^{j-1})$  and hence  $\mathcal{R}(K^{-1}[Z_k, z]) \subseteq \mathcal{R}(S^j)$ . Consequently,  $(I - Y_k H_k^{-1}[Q_k, q]^*)r \in \mathcal{R}(S^j)$ . □

**Lemma 4.4.** Let  $r \in \mathcal{R}(S^j)$ ,  $K = A - \tau_0 B$ ,  $AQ_k = Z_k S_A$ ,  $BQ_k = Z_k S_B$ ,  $q \in \mathcal{R}(S^j)$ ,  $z = \nu Aq + \mu Bq$ ,  $Y_k = K^{-1}[Z_k, z]$  and  $H_k = [Q_k, q]^* Z_k$ . Then

$$\mathcal{K}^j((I - Y_k H_k^{-1} Q_k^*) K^{-1} (\beta A - \alpha B) (I - Y_k H_k^{-1} Q_k^*), r) \subseteq \mathcal{R}(S^j).$$

*Proof.* The result follows from applying subsequently lemma 4.3, 4.1, 4.2 and again 4.3. □

This lemma not only enables one to use a Krylov solver for the correction equation, it also has consequences for purification in Jacobi-Davidson.

**Lemma 4.5.** If the initial search space  $V_0 \subset \mathcal{R}(S^j)$  and the Jacobi-Davidson correction equation is solved using an exact preconditioner, then all subsequent search spaces  $V_k \subset \mathcal{R}(S^j)$ .

*Proof.* The result follows from lemma 4.4. □

**Theorem 4.6.** If the initial search space  $V_0 \subset \mathcal{R}(S^j)$  and the Jacobi-Davidson correction equation is solved using an exact preconditioner, then in exact arithmetic no spurious eigenpairs are computed during the Jacobi-Davidson process.

*Proof.* The reduced system is  $(M_A, M_B) = (W^* A V, W^* B V)$  with test space  $W = \nu A V + \mu B V$ . Applying lemma 4.1 to  $W$  gives  $W \subset \mathcal{R}(BS^{j-1})$  and no spurious eigenvalues are computed. From lemma 4.5 it follows that the Petrov vectors  $q_i = V_k s_i$  satisfy  $q_i \in \mathcal{R}(S^j)$ . □

The last theorem says that, in exact arithmetic, if the Jacobi-Davidson method with exact preconditioning starts with  $V_0 \subset \mathcal{R}(S^j)$ , then  $V_k \subset \mathcal{R}(S^j)$  and  $W \subset \mathcal{R}(BS^{j-1})$  and no spurious eigenpairs are computed. In other words, with exact preconditioning the search space is purified automatically. The effect is even enforced because usually more than one iteration of the Krylov solver is needed.

However, in finite arithmetic components in  $\mathcal{N} + \mathcal{G}$  may still arise due to rounding errors, and if an exact preconditioner is not available, theorem 4.6 is also not applicable. Fortunately, there is a result similar to theorem 3.5. Let  $P_{\mathcal{R}}$ ,  $P_{\mathcal{N}}$  and  $P_{\mathcal{G}}$  be normalized projectors that map a vector into  $\mathcal{R} = \mathcal{R}(A^{-1}B)$ ,  $\mathcal{N} = \mathcal{N}(A^{-1}B)$  and  $\mathcal{G} = \mathcal{G}(A^{-1}B)$  respectively, so  $x \in \mathbb{C}^m$  can be decomposed uniquely as  $x = P_{\mathcal{R}}x + P_{\mathcal{N}}x + P_{\mathcal{G}}x$ . The following theorem shows that a converged Petrov pair  $(\lambda, x)$  is purified automatically, provided  $|\lambda|$ ,  $\|A^{-1}\|_2$  and  $\|B\|_2$  are not too large.

**Theorem 4.7.** Let  $(\lambda, x)$  be a converged Petrov pair of  $(A, B)$ , with  $r = Ax - \lambda Bx$  and  $\|r\|_2 < \epsilon$ . Then

$$\begin{aligned} \|P_{\mathcal{N}}x\|_2 &\leq \epsilon \|A^{-1}\|_2 (1 + |\lambda| \|A^{-1}\|_2 \|B\|_2), \\ \|P_{\mathcal{G}}x\|_2 &\leq \epsilon \|A^{-1}\|_2. \end{aligned}$$

*Proof.* Use  $x = A^{-1}(r + \lambda Bx)$ ,  $P_{\mathcal{N}}(A^{-1}B) = (A^{-1}B)P_{\mathcal{G}}$  and  $P_{\mathcal{G}}(A^{-1}B) = 0$ . □

#### 4.4 Harmonic Ritz-values, exact targets and purification

Numerical experiments show that the JDQZ process with harmonic Petrov values and a target equal to an eigenvalue does not converge to this eigenvalue, but to eigenvalues closest to the target. This observation can be understood from a theoretical point of view, as will be explained next, and may be of use in avoiding convergence to eigenvalues at infinity.

First consider the Jacobi-Davidson process for the ordinary eigenproblem

$$Ax = \lambda x$$

In [20] it is shown that the harmonic Ritz values of  $A$  are equal to the eigenvalues of the  $k \times k$  matrix

$$\tilde{H}_k = (W_k^* V_k)^{-1} W_k^* A V_k = (W_k^* V_k)^{-1}$$

with  $W_k = AV_k$  and  $W_k^* W_k = I$ . Note that  $\tilde{H}_k^{-1} = W_k^* V_k = W_k^* A^{-1} W_k$ , the projection of  $A^{-1}$  with respect to an orthonormal basis  $W_k$ . In practice it is not necessary to invert  $\tilde{H}_k^{-1}$ , because the harmonic Ritz values of  $A$  are the reciprocals of the eigenvalues of  $\tilde{H}_k = W_k^* V_k$ . Hence, no problems are encountered if  $W_k^* V_k$  is singular, which may happen if  $A$  has an eigenvalue at zero.

**Theorem 4.8.** *Let  $A \in \mathbb{R}^{n \times n}$  be a normal matrix and  $\tau \in \mathbb{R}$ . If  $\tau$  exactly equals an eigenvalue of  $A$ , then, in exact arithmetic, the Jacobi-Davidson process with harmonic Ritz values and target  $\tau$  will not converge to the eigenvalue  $\lambda = \tau$ .*

*Proof.* Without loss of generality, let  $\tau = \lambda = 0$ : if  $\tau = \lambda \neq 0$ , the proof follows for  $A - \tau I$ . Denote the null space of  $A$  by  $\mathcal{N}$  and the range of  $A$  by  $\mathcal{R}$ . The eigenspace corresponding to the eigenvalue  $\lambda = 0$  is (a subset of) the null space  $\mathcal{N}$ . However, because of the normality of  $A$ , the space spanned by the columns of  $W_k = AV_k$  does not contain any elements of the eigenspace of  $\lambda = 0$ . Because the eigenspaces of  $A$  and  $A^{-1}$  are the same, the proof would be complete if  $A^{-1}$  would exist. However, in this case there is an eigenvalue  $\lambda = 0$  and hence  $A^{-1}$  does not exist.

If  $\delta \in \mathbb{C} \setminus \Lambda(A)$ , then  $(A + \delta I)^{-1}$  exists and the eigenspaces of  $A$ ,  $A + \delta I$ , and  $(A + \delta I)^{-1}$  are the same, but the eigenvalues are  $\lambda$ ,  $\lambda + \delta$ , and  $(\lambda + \delta)^{-1}$  respectively. With the iteration vectors  $w_k$  still generated by  $Av_k$ , and hence  $W_k$  containing no components of the null space of  $A$  and the eigenspace  $V_\delta$  of  $A + \delta I$ , it follows that the eigenvalue  $\delta^{-1}$  will not be contained in the set of eigenvalues of  $W_k^* (A + \delta I)^{-1} W_k$ .  $\square$

In finite arithmetic,  $W_k$  can still contain components of the (generalized) null space of  $A$ , which may hamper convergence to the desired eigenvalues or even cause convergence to the undesired, perturbed eigenvalue. If the starting vector is in the null space of  $A$ , Jacobi-Davidson with harmonic Ritz values will break down.

The proof for the Jacobi-Davidson  $QZ$  process for generalized eigenproblems is similar. In [21] these observations are used to derive a selection strategy for Ritz pairs.

#### 4.5 A strategy with JDQZ

The strategy in Alg. 5 is conceptually the same as Alg. 4, with JDQZ instead of IRA. By considering the pencil  $(B, A)$  instead of  $(A, B)$ , the infinite eigenvalues are transformed to zero. The extremal eigenvalues of  $(B, A)$  can be computed safely, efficiently and accurately by JDQZ with harmonic Petrov values and target  $\tau = 0$  (see theorem 4.8). These eigenvalues can be used to determine  $\alpha_1, \alpha_2$  and  $\alpha_3 = 0$  for the modified Cayley transform (see also section 3.3), here formulated as the generalized eigenvalue problem  $\mathcal{A}(\alpha_2, \alpha_3)x = \mu \mathcal{B}(\alpha_1)x$  with

$$\mathcal{A}(\alpha_2, \alpha_3) = \begin{bmatrix} K - \alpha_2 M & \alpha_3 C \\ \alpha_3 C^T & 0 \end{bmatrix}, \quad \mathcal{B}(\alpha_1) = \begin{bmatrix} K - \alpha_1 M & C \\ C^T & 0 \end{bmatrix}.$$

Eigenpairs that are found in phase 1 (step 1 - 3) can be deflated from the problem in phase 2 (step 4 - 6).



---

**Algorithm 5** Strategy for computing the 2 left most eigenvalues of  $(A, B)$ 

---

**PHASE 1**

- 1: Choose a suitable preconditioner for the correction equation
- 2: Compute the  $r \geq 2$  largest eigenvalues  $\tilde{\lambda}_i$  of  $(B, A)$  with JDQZ ( $\tau = 0$ )
- 3: Order  $\lambda_i = 1/\tilde{\lambda}_i$ ,  $i = 1, \dots, r$  by increasing real part

**PHASE 2**

- 4: Determine  $\alpha_1$  and  $\alpha_2$ , and  $\alpha_3 = 0$
  - 5: Compute the largest 2 eigenvalues  $\mu_i$  of  $(\mathcal{A}(\alpha_2, \alpha_3), \mathcal{B}(\alpha_1))$  with JDQZ
  - 6: The eigenvalues of  $(A, B)$  are  $\lambda_i = \frac{\alpha_1 \mu_i - \alpha_2}{\mu_i - 1}$
- 

## 4.6 Realistic examples

The strategy in Alg. 5 is applied to the test problems of section 3.4. To make a fair comparison with the IRA strategy in Alg. 4, two situations were considered:

- The correction equation is not solved exactly, but with 20 steps of (unrestarted) GMRES [17] with preconditioner  $A$  (stopping earlier if the relative residual norm drops below  $\epsilon = 10^{-6}$ ).
- The correction equation is solved exactly and an initial search space of size  $j_{\min}$  is computed with Arnoldi.

In both situations, the initial vector  $v_1$  had all entries one and was not purified. The search and test space dimensions are limited by  $j_{\min} = 15$  and  $j_{\max} = 20$ , and the residual tolerance was  $10^{-6}$  (see [7] for more details about the several parameters and sophisticated stopping criteria). In situation 1, solves with preconditioner  $A$  are needed and hence one could argue that in that case also the IRA strategy in Alg. 4 could be used. The goal here however is to show that even if the correction equation is not solved exactly and the initial search space is not constructed with Arnoldi, JDQZ is able to compute the leftmost eigenvalues. In this way, situation 1 resembles the situation where indeed solves with  $A$  are not possible and the IRA strategy is not applicable. The quality of the preconditioner influences the speed of convergence of the GMRES process, but this paper is not concerned with designing a good preconditioner (see [4, chapter 8] and references therein for preconditioners of related systems). For the experiments, a variant of the JDQZ algorithm, that keeps the search and test spaces real, is used (RJDQZ [24]).

### 4.6.1 Flow over a backward facing step

Table 3 shows statistics for Alg. 5 with  $r = 2$ , for both exact and inexact solution of the correction equation. The leftmost eigenvalues  $\lambda_1 = 6.04 \cdot 10^{-2}$  and  $\lambda_{2,3} = 7.97 \cdot 10^{-2} \pm i1.92 \cdot 10^{-2}$  were already found in the first phase of the algorithm: the validation in phase 2 did not result in new eigenvalues. The differences in residual norms can be explained by the asymptotically quadratical convergence of the exact variant. Concerning the higher computing times for the inexact variant, one should keep in mind that for stability analysis the quality of the solution (no missed eigenvalues) is the most important. Furthermore, it may be expected that the times can be decreased by using a more effective preconditioning, but this goes beyond the scope of this paper. The exact variant is faster than implicitly restarted Arnoldi (cf. table 2).

### 4.6.2 Flow in a driven cavity

Table 4 shows statistics for Alg. 5 with  $r = 2$ , for both exact and inexact solution of the correction equation. The eigenvalues  $\lambda_1 = 3.21 \cdot 10^{-2}$  and  $\lambda_4 = 1.01 \cdot 10^{-1}$  were found in the first phase of the algorithm. The validation in phase 2 found the missed eigenvalue  $\lambda_{2,3} = 6.20 \cdot 10^{-2} \pm i4.61 \cdot 10^{-1}$ . The exact variant is faster than implicitly restarted Arnoldi (cf. table 2).

Table 3: Statistics for Alg. 5 for the backward facing step with Reynolds number  $Re = 800$  (section 4.6.1): number of iterations, restarts, time, found eigenvalues and residuals after each phase. No restarts were needed.

	Inexact		Exact	
	phase 1	phase 2	phase 1	phase 2
#iterations	18	9	4	2
time (s)	1400	1000	85	80
eigenvalues	$\lambda_1, \lambda_{2,3}$	$\lambda_1$	$\lambda_1, \lambda_{2,3}$	$\lambda_1, \lambda_{2,3}$
$\max_i \ Ax_i - \lambda_i Bx_i\ $	$1 \cdot 10^{-11}$	$1 \cdot 10^{-11}$	$1 \cdot 10^{-15}$	$1 \cdot 10^{-15}$

Table 4: Statistics for Alg. 4 for the driven cavity with Reynolds number  $Re = 500$  (section 4.6.2): number of restarts, time, found eigenvalues and residuals after each phase. No restarts were needed.

	Inexact		Exact	
	phase 1	phase 2	phase 1	phase 2
#iterations	13	12	3	3
time (s)	330	340	21	58
eigenvalues	$\lambda_1, \lambda_4$	$\lambda_1, \lambda_{2,3}$	$\lambda_1, \lambda_4$	$\lambda_1, \lambda_{2,3}$
$\max_i \ Ax_i - \lambda_i Bx_i\ $	$1 \cdot 10^{-10}$	$1 \cdot 10^{-10}$	$1 \cdot 10^{-15}$	$1 \cdot 10^{-15}$

## 5 Conclusions

The strategy based on implicitly restarted Arnoldi is a reliable and fast method to compute the leftmost eigenvalues of large scale eigenvalues, if the solves needed for the shift-and-invert and Cayley transformations can be done efficiently and exactly. By exploiting the structure of the generalized eigenvalue problem and by choosing suitable parameters for the modified Cayley transformation, the troubles caused by infinite eigenvalues are circumvented and no purification is needed.

If the solves that are needed for the transformations cannot be done exactly, the Jacobi-Davidson QZ method is a good alternative. Following the same strategy, JDQZ is able to compute the leftmost eigenvalues, without corruption due to infinite eigenvalues. If solves can be done exactly, it is faster than implicitly restarted Arnoldi. Jacobi-Davidson puts no requirements on the matrix pencil: it can handle both regular and singular pencils.

## Acknowledgments

I thank Tycho van Noorden for many fruitful discussions. The result in lemma 4.5 was independently derived by both authors. Tycho van Noorden uses the result in a continuation method context. Gerard Sleijpen pointed me to [14] and gave helpful comments. I thank Henk van der Vorst for useful comments on earlier versions of this text. I am thankful to David Silvester and Howard Elman for introducing me to IFISS [18]. I am grateful to the anonymous referee, whose suggestions and remarks helped me to improve the paper.

## References

- [1] K. A. Cliffe, T. J. Garratt, and A. Spence, *Eigenvalues of the discretized Navier-Stokes equation with application to the detection of Hopf bifurcations*, Adv. Math. Comp. **1** (1993), 337–356.

- [2] ———, *Eigenvalues of block matrices arising from problems in fluid mechanics*, SIAM J. Matrix Anal. Appl. **15** (1994), no. 4, 1310–1318.
- [3] K. A. Cliffe, A. Spence, and S. J. Tavener, *The numerical analysis of bifurcation problems with application to fluid mechanics*, Acta Numerica **9** (2000), 39–131.
- [4] H. Elman, D. Silvester, and A. Wathen, *Finite Elements and Fast Iterative Solvers*, first ed., Oxford University Press, 2005.
- [5] T. Ericsson, *A generalised eigenvalue problem and the Lanczos algorithm*, Large Scale Eigenvalue Problems (J. Cullum and R.A. Willoughby, eds.), Elsevier Science Publishers BV, 1986, pp. 95–119.
- [6] T. Ericsson and A. Ruhe, *The spectral transformation lanczos method for the numerical solution of large sparse generalized symmetric eigenvalue problems*, Math. Comp. **35** (1980), 1251–1268.
- [7] D. R. Fokkema, G. L. G. Sleijpen, and H. A. van der Vorst, *Jacobi-Davidson style QR and QZ algorithms for the reduction of matrix pencils*, SIAM J. Sc. Comp. **20** (1998), no. 1, 94–125.
- [8] T. J. Garratt, *The numerical detection of Hopf bifurcations in large systems arising in fluid mechanics*, Ph.D. thesis, University of Bath, 1991.
- [9] P. M. Gresho, D. K. Gartling, J. R. Torczynski, K. A. Cliffe, K. H. Winters, T. J. Garratt, A. Spence, and J. W. Goodrich, *Is the steady viscous incompressible two-dimensional flow over a backward-facing step at  $Re=800$  stable?*, Int. J. Num. Meth.Fluids **17** (1993), 501–541.
- [10] R. B. Lehoucq and J. A. Scott, *Implicitly restarted Arnoldi methods and eigenvalues of the discretized Navier Stokes equations*, Tech. Report SAND97-2712J, Sandia National Laboratory, 1997.
- [11] R. B. Lehoucq and D. C. Sorensen, *Deflation techniques within an implicitly restarted Arnoldi iteration*, SIAM J. Matrix Anal. Appl. **17** (1996), 789–821.
- [12] R. H. Lehoucq, D. C. Sorensen, and C. Yang, *ARPACK SOFTWARE*, <http://www.caam.rice.edu/software/ARPACK/>.
- [13] K. Meerbergen and D. Roose, *The restarted Arnoldi method applied to iterative linear system solvers for the computation of rightmost eigenvalues*, SIAM J. Matrix Anal. Appl. **18** (1997), no. 1, 1–20.
- [14] K. Meerbergen and A. Spence, *Implicitly restarted Arnoldi with purification for the shift-invert transformation*, Math. Comp. **66** (1997), no. 218, 667–689.
- [15] K. Meerbergen, A. Spence, and D. Roose, *Shift-invert and Cayley transforms for detection of rightmost eigenvalues of nonsymmetric matrices*, BIT **34** (1994), no. 3, 409–423.
- [16] B. Nour-Omid, B. N. Parlett, T. Ericsson, and P. S. Jensen, *How to implement the spectral transformation*, Math. Comp. **48** (1987), no. 178, 663–673.
- [17] Y. Saad and M. H. Schultz, *GMRES: A generalized minimal residual algorithm for solving nonsymmetric linear systems*, SIAM J. Sci. Stat. Comput. **7** (1986), no. 3, 856–869.
- [18] D. Silvester, H. Elman, and A. Ramage, *Incompressible Flow and Iterative Solver Software*, <http://www.maths.manchester.ac.uk/djs/ifiss/>.
- [19] G. L. G. Sleijpen, J. G. L. Booten, D. R. Fokkema, and H. A. van der Vorst, *Jacobi-Davidson type methods for generalized eigenproblems and polynomial eigenproblems*, BIT **36** (1996), no. 3, 595–633.

- [20] G. L. G. Sleijpen and H. A. van der Vorst, *A Jacobi-Davidson iteration method for linear eigenvalue problems*, SIAM J. Matrix Anal. Appl. **17** (1996), no. 2, 401–425.
- [21] G. L. G. Sleijpen, H. A. van der Vorst, and E. Meijerink, *Efficient expansion of subspaces in the Jacobi-Davidson method*, Electronic Transactions on Numerical Analysis **7** (1998), 75–89.
- [22] D. C. Sorensen, *Implicit application of polynomial filters in a  $k$ -step Arnoldi method*, SIAM J. Matrix Anal. Appl. **13** (1992), 357–385.
- [23] ———, *Deflation for implicitly restarted Arnoldi methods*, Tech. Report TR98-12, Rice University, 1998.
- [24] T. L. van Noorden and J. Rommes, *Computing a partial real ordered generalized Schur form using the Jacobi-Davidson method*, RANA 05-41, Eindhoven Technical University, 2005, Accepted for publication in Num. Lin. Alg. Appl.