

A NOTE ON THE GREEDY β -TRANSFORMATION WITH DELETED DIGITS

KARMA DAJANI AND CHARLENE KALLE

ABSTRACT. In this article we give the attractor and the absolutely continuous, invariant measure of the greedy and lazy β -transformation with deleted digits and show their ergodicity. We will consider two specific examples of greedy β -transformations of which the invariant measure can be explicitly calculated.

1. INTRODUCTION

Invariant measures for β -transformations have been an intensively studied subject for years. The articles by A. Rényi ([12]) and W. Parry ([9]) are fundamental in this respect. Both consider the greedy transformation for a $\beta > 1$ and a digit set $\{0, 1, \dots, \lfloor \beta \rfloor\}$ on the unit interval. This transformation $T_\beta : [0, 1] \rightarrow [0, 1]$ is given by $T_\beta x = \beta x \pmod{1}$. A. Rényi proved the existence of a unique invariant measure, equivalent to the Lebesgue measure, for the greedy β -transformation and W. Parry gave an explicit expression for the density function of this measure. In [5] F. Hofbauer proved that the measure obtained by W. Parry is the unique measure of maximal entropy.

A number of articles have been published on invariant measures of piecewise monotonic transformations. Among others, the articles [1] by J. Buzzi and O. Sarig, [2] by W. Byers and A. Boyarsky, [3] by W. Byers, P. Góra and A. Boyarsky, [5] and [6] by F. Hofbauer, [7] by A. Lasota and J. Yorke, [8] by T.Y. Li and J. Yorke, [13] by F. Schweiger and [14] by K. Wilkinson state a variety of results regarding invariant measures of this kind of transformations and their ergodicity.

In [4] a definition is given of the greedy and lazy β -transformations with deleted digits, which are generalizations of the classical greedy transformation mentioned earlier. The definition in that paper is a dynamical one and is based on an algorithm, given by M. Pedicini in [11]. For each $\beta > 1$ and digit set $A = \{a_0, \dots, a_m\}$ with $a_0 < \dots < a_m$, that satisfies the following condition

$$(1) \quad \max_{0 \leq j \leq m-1} (a_{j+1} - a_j) \leq \frac{a_m - a_0}{\beta - 1},$$

the greedy transformation with deleted digits,

$$T = T_{\beta, A} : \left[\frac{a_0}{\beta - 1}, \frac{a_m}{\beta - 1} \right] \rightarrow \left[\frac{a_0}{\beta - 1}, \frac{a_m}{\beta - 1} \right]$$

1991 *Mathematics Subject Classification.* Primary, 37A05, 11K55.

Key words and phrases. greedy and lazy expansions, attractor, invariant measure.

can be defined as follows.

$$Tx = \begin{cases} \beta x - a_j, & \text{if } x \in \left[\frac{a_0}{\beta-1} + \frac{a_j - a_0}{\beta}, \frac{a_0}{\beta-1} + \frac{a_{j+1} - a_0}{\beta} \right), \\ & \text{for } j = 0, \dots, m-1, \\ \beta x - a_m, & \text{if } x \in \left[\frac{a_0}{\beta-1} + \frac{a_m - a_0}{\beta}, \frac{a_m}{\beta-1} \right]. \end{cases}$$

A set $A = \{a_0, \dots, a_m\}$ with $a_0 < \dots < a_m$ that satisfies condition (1) for a $\beta > 1$ is called an *allowable digit set for β* . The lazy transformation with deleted digits is defined in [4] in a similar way. For each $\beta > 1$ and each allowable digit set $A = \{a_0, \dots, a_m\}$, the lazy transformation with deleted digits,

$$L = L_{\beta, A} : \left[\frac{a_0}{\beta-1}, \frac{a_m}{\beta-1} \right] \rightarrow \left[\frac{a_0}{\beta-1}, \frac{a_m}{\beta-1} \right]$$

is given by

$$Lx = \begin{cases} \beta x - a_0, & \text{if } x \in \left[\frac{a_0}{\beta-1}, \frac{a_m}{\beta-1} - \frac{a_m - a_0}{\beta} \right], \\ \beta x - a_j, & \text{if } x \in \left(\frac{a_m}{\beta-1} - \frac{a_m - a_{j-1}}{\beta}, \frac{a_m}{\beta-1} - \frac{a_m - a_j}{\beta} \right], \\ & \text{for } j = 1, \dots, m. \end{cases}$$

The algorithm by M. Pedicini that can be found in [11] generates the same β -expansions as the greedy β -transformation with deleted digits. In [11], M. Pedicini showed that this algorithm is well-defined if the digit set A satisfies condition (1), and in [4] the same was shown for the greedy and lazy transformation. The difference with the classical versions of these two transformations is the use of a more general digit set.

In this article we will study the greedy transformation with deleted digits with respect to its invariant measure that is absolutely continuous with respect to the Lebesgue measure. In [4] it is shown that for each $\beta > 1$ and each allowable digit set $A = \{a_0, \dots, a_m\}$ the greedy transformation with deleted digits $T_{\beta, A}$ is isomorphic to the greedy transformation T_{β, A^*} , where A^* is an allowable digit set for β , containing $m+1$ digits, but for which $a_0^* = 0$. So without loss of generality we assume that $a_0 = 0$ for each allowable digit set A . It was also shown in the same article that the greedy transformation with deleted digits T given for $\beta > 1$ and digit set $A = \{a_0, \dots, a_m\}$ is isomorphic to the lazy transformation with deleted digits L , defined for the same $\beta > 1$, but with digit set $\tilde{A} = \{\tilde{a}_0, \dots, \tilde{a}_m\}$, where $\tilde{a}_i = a_0 + a_m - a_{m-i}$, $i \in \{0, \dots, m\}$. This isomorphism ϕ is given by

$$\phi : \left[\frac{a_0}{\beta-1}, \frac{a_m}{\beta-1} \right] \rightarrow \left[\frac{\tilde{a}_0}{\beta-1}, \frac{\tilde{a}_m}{\beta-1} \right] : x \mapsto \frac{a_0 + a_m}{\beta-1} - x.$$

So we have $L \circ \phi = \phi \circ T$. Notice that if $a_0 = 0$, then also $\tilde{a}_0 = 0$, so without loss of generality we can assume that the lazy transformation L is defined for a $\beta > 1$ and an allowable digit set $\tilde{A} = \{\tilde{a}_0, \dots, \tilde{a}_m\}$, such that $\tilde{a}_0 = 0$. Notice that then $\tilde{a}_m = a_m - a_0 = a_m$, so that we have $\tilde{a}_i = a_m - a_{m-i}$ and $a_i = \tilde{a}_m - \tilde{a}_{m-i}$.

In the first section of this article we will prove the existence of a unique absolutely continuous, invariant measure for the greedy transformation with deleted digits using the results found by T.Y. Li and J. Yorke in [8], and give its support. We will

make similar observations for the lazy transformation with deleted digits and we will also make a remark about the ergodicity of the greedy and lazy transformation. In the last section we consider in more detail two classes of greedy transformations with deleted digits, and we give an explicit formula for the density of their absolutely continuous invariant measures. For the first class an article by K. Wilkinson ([14]) has been an important source. For the second class we use an article by W. Byers and A. Boyarski [2]), which is based on [10] by W. Parry.

2. ATTRACTOR

Let $\beta > 1$ and $A = \{a_0, a_1, \dots, a_m\}$ be an allowable digit set with $a_0 = 0$. Let $T = T_{\beta, A} : \left[0, \frac{a_m}{\beta - 1}\right] \rightarrow \left[0, \frac{a_m}{\beta - 1}\right]$ be the greedy transformation with deleted digits. This is a piecewise linear, strictly increasing transformation, which has its discontinuities in the points $\frac{a_i}{\beta}$ for $i = 1, \dots, m$. Let J denote the set containing these points. Then J is finite and for each $x \in \left[0, \frac{a_m}{\beta - 1}\right] \setminus J$ we have $T'(x) = \beta > 1$. The points in J give a partition $\Delta = \{\Delta_i\}_{i=0}^m$ of the interval $\left[0, \frac{a_m}{\beta - 1}\right]$, where $\Delta_m = \left[\frac{a_m}{\beta}, \frac{a_m}{\beta - 1}\right]$ and for $i \in \{0, \dots, m - 1\}$, $\Delta_i = \left[\frac{a_i}{\beta}, \frac{a_{i+1}}{\beta}\right)$. Define for $i \in \{1, \dots, m\}$ the values y_i to be the values obtained from T by taking the limit from the left in the points $\frac{a_i}{\beta}$, i.e. $y_i = a_i - a_{i-1}$. (See Figure 1)

We will begin by defining different notions of invariance under the transformation T and by stating the results found by A. Lasota and J. Yorke in [7] and by T.Y. Li and J. Yorke in [8]. A measure μ is called *an invariant measure under T* if for all measurable sets E we have $\mu(E) = \mu(T^{-1}E)$. A μ -integrable function f is called *an invariant function under T* if for all measurable sets E , $\int_E f d\mu = \int_{T^{-1}E} f d\mu$. And we will call a measurable set E *forward invariant under T* if TE is measurable and $TE = E$ modulo sets of measure zero. Let λ denote the Lebesgue measure. It was shown in [7] that there exists an invariant measure, absolutely continuous with respect to the Lebesgue measure, for transformations τ that are piecewise continuous with a finite set of points of discontinuity and that have a derivative bigger than 1 for points outside this finite set. In [8], Li and Yorke studied these invariant measures in more detail and their results translate in the following way to our particular greedy transformation T . For the transformation T , there exist sets B_1, \dots, B_n and functions f_1, \dots, f_n , where $n \leq m$, such that all the following hold.

- (c1) For each $i \in \{1, \dots, n\}$, B_i is a finite union of closed subintervals of $\left[0, \frac{a_m}{\beta - 1}\right]$. Each B_i contains at least one of the elements of J in its interior. Moreover, each B_i is forward invariant under T .
- (c2) $B_i \cap B_j$ contains at most a finite number of points, when $i \neq j$, $i, j \in \{1, \dots, n\}$.




FIGURE 1. The greedy β -transformation with $\beta = 2.5$ and $A = \{0, 1.35, 1.75, 3.3, 6\}$.

- (c3) For almost all $x \in \left[0, \frac{a_m}{\beta-1}\right] \setminus J$, there is an $i \in \{1, \dots, n\}$ such that the closure of the forward orbit of x under T equals the set B_i , i.e.

$$\Lambda(x) := \bigcap_{N=1}^{\infty} \overline{\{T^n(x)\}_{n=N}^{\infty}} = B_i.$$

- (c4) For each $i \in \{1, \dots, n\}$, B_i is the support of the function f_i , i.e. $f_i > 0$ λ a.e. on B_i and $f_i = 0$ on B_i^c . Moreover, $\int_{B_i} f_i d\lambda = 1$.
- (c5) For each $i \in \{1, \dots, n\}$, f_i is invariant under T and if a function g satisfies (c4) for some i , then $g = f_i$ λ almost everywhere.

- (c6) Each function f that is invariant under T can be written as $f = \sum_{i=1}^n b_i f_i$ with a suitably chosen set of constants $\{b_i\}_{i=1}^n$.

- (c7) If f is an invariant function and E is a measurable set, such that TE is measurable and $TE \subseteq E$ λ a.e., then $f \cdot 1_E$ is an invariant function, where 1_E denotes the indicator function of the set E .

Remark 2.1. The last result was proven in [8] for sets E such that $TE = E$ for λ a.e. $x \in E$, but the proof of T.Y. Li and J. Yorke still holds under the weaker assumption that $TE \subseteq E$ λ a.e.

We will first make some observations about the sets B_i .

Lemma 2.1. *Let $I \subseteq \left[0, \frac{a_m}{\beta - 1}\right]$ be a closed interval.*

(i) *If I is forward invariant under T and contains at least one element of J in its interior, then $0 \in I$.*

(ii) *If I does not contain an element of J in its interior, then I is not forward invariant under T .*

Proof. The first part of the lemma follows immediately from the fact that for each $i \in \{1, \dots, m\}$, $T\left(\frac{a_i}{\beta}\right) = 0$.

For the second part it is enough to notice that if I does not contain an element of J in its interior, then $\lambda(TI) = \beta\lambda(I)$. \square

Remark 2.2. As an immediate consequence of this lemma, we have that there cannot exist two or more sets satisfying (c1) and (c2). To see this, suppose that the sets B_i and B_j both satisfy (c1). Then they are both forward invariant under T , so by the previous lemma there should exist numbers $0 < x_i, x_j \leq \frac{a_m}{\beta - 1}$ such that $[0, x_i] \subseteq B_i$ and $[0, x_j] \subseteq B_j$, but this contradicts (c2). So by the previously stated results from [8], we know that there exists a number $0 < x \leq \frac{a_m}{\beta - 1}$ and a

finite number of closed intervals $I_1, \dots, I_k \subseteq \left[0, \frac{a_m}{\beta - 1}\right]$ such that the set

$$(2) \quad B := [0, x] \cup \bigcup_{i=1}^k I_i$$

satisfies (c1) to (c7) for an invariant probability density function f . This means that there exists a unique invariant measure for T that is absolutely continuous with respect to the Lebesgue measure on B . Notice that without loss of generality we can assume that B is the finite union of disjoint, closed intervals. The fact that B is forward invariant then implies that $T[0, x] \subseteq [0, x]$.

The next lemma states that if B contains a closed interval whose image under T is contained in itself, then B is exactly this interval.

Lemma 2.2. *Let f be the invariant probability density as in Remark 2.2 and let B be its support. Suppose that $[\alpha_1, \alpha_2] \subseteq B$ is a closed interval. If $T[\alpha_1, \alpha_2] \subseteq [\alpha_1, \alpha_2]$ λ a.e., then $[\alpha_1, \alpha_2] = B$. Consequently, $[\alpha_1, \alpha_2]$ is a forward invariant set.*

Proof. Consider the function $g = f \cdot 1_{[\alpha_1, \alpha_2]}$. Since f is an invariant function and $[\alpha_1, \alpha_2]$ satisfies $T[\alpha_1, \alpha_2] \subseteq [\alpha_1, \alpha_2]$ λ a.e., by (c7) we know that also the function g is invariant, with its support contained in the support of f . By (c6) this means that there exists a constant c , such that $g = c \cdot f$. Now define the function

$$h = \frac{g}{\int g d\lambda}.$$

Then h is an invariant probability density function and $h = c' \cdot f$, with $c' = c / \int g d\lambda$. This means that $h = f$ λ a.e., so that $1_{[\alpha_1, \alpha_2]}(x) = 1$ for λ almost all $x \in B$. Since B is a finite union of closed intervals, it follows that $B = [\alpha_1, \alpha_2]$. By (c1), $[\alpha_1, \alpha_2]$ is forward invariant. \square

Remark 2.3. By the same reasoning as in the proof of the previous lemma, it can be shown that T is ergodic with respect to the invariant measure. To see this, let μ be the measure given by $\mu(E) = \int_E f d\lambda$ for each measurable set E and suppose that A is a measurable set such that $T^{-1}A = A$ λ a.e. and $\mu(A) > 0$. Then $TA \subseteq A$ λ a.e., so by (c7) the function $g = f \cdot 1_A$ is invariant. Following the idea of the proof of Lemma 2.2 gives that $1_A = 1$ λ a.e., so $\mu(A) = 1$.

By Remark 2.2 we know that there exists an element $x \in \left[0, \frac{a_m}{\beta - 1}\right]$, such that the support B of f contains the interval $[0, x]$ and $T[0, x] \subseteq [0, x]$. By Lemma 2.2, this means that $B = [0, x] = T[0, x]$ λ a.e. The next two lemmas specify the value of x . First we define the following value. Let $y_{i_0} = \max \left\{ y_i : \frac{a_i}{\beta} \leq x \right\}$ and if there are two such values, then let y_{i_0} be the one with the smallest index.

Lemma 2.3. *Let $B = [0, x]$ be the support of the probability density function f , as described above. Then $B = [0, y_{i_0}]$.*

Proof. Since $T[0, x] = [0, x]$ λ a.e., we have that $y_i \leq x$ for any i such that $\frac{a_i}{\beta} \leq x$. Hence $y_{i_0} \leq x$. Also $Tx \leq x$.

Suppose $x \in \Delta_k$ for some $k \in \{0, \dots, m\}$. Then by the definition of y_{i_0} ,

$$(3) \quad T \left[0, \frac{a_k}{\beta} \right) \subseteq [0, y_{i_0}] \quad \lambda \text{ a.e.}$$

If $y_{i_0} \in \left[0, \frac{a_k}{\beta}\right]$, then $T[0, y_{i_0}] \subseteq [0, y_{i_0}] \subseteq [0, x]$ λ a.e. and thus by Lemma 2.2, $[0, y_{i_0}] = [0, x]$ λ a.e. If on the other hand $y_{i_0} \in \left[\frac{a_k}{\beta}, x\right]$, then since $Tx \leq x$, we also have $Ty_{i_0} \leq y_{i_0}$ and this means that

$$T \left[\frac{a_k}{\beta}, y_{i_0} \right] \subseteq [0, y_{i_0}] \quad \lambda \text{ a.e.}$$

Combining this with equation (3) gives that $T[0, y_{i_0}] \subseteq [0, y_{i_0}] \subseteq [0, x]$ λ a.e., so again by Lemma 2.2 we have that $[0, y_{i_0}] = [0, x]$. \square

From the previous lemma we know that x is one of the values y_i , $i \in \{1, \dots, m\}$. The next lemma states explicitly which of these values it is.

Lemma 2.4. *Let y_{i_0} be defined as above. Then*

$$(4) \quad i_0 = \min \{ i : T[0, y_i] \subseteq [0, y_i] \quad \lambda \text{ a.e.} \}.$$

Proof. Since $[0, x] = [0, y_{i_0}]$ is the support of the invariant probability density function f , we must have by Lemma 2.2 that $T[0, y_i] \not\subseteq [0, y_i]$ λ a.e. for any $y_i < y_{i_0}$. In particular, by the definition of y_{i_0} we have that if $i < i_0$, then

$$\frac{a_i}{\beta} < \frac{a_{i_0}}{\beta} \leq y_{i_0}.$$

This implies that $y_i < y_{i_0}$ and thus that $T[0, y_i] \not\subseteq [0, y_i] \lambda$ a.e. Hence $i_0 = \min\{i : T[0, y_i] \subseteq [0, y_i] \lambda \text{ a.e.}\}$. \square

In the previous lemmas and remarks we have established the existence of a unique absolutely continuous, invariant measure for the greedy transformation with deleted digits, that is ergodic, and we have given its support. These results are summarized in the following theorem.

Theorem 2.1. *Let $\beta > 1$ and let $A = \{0, a_1, \dots, a_m\}$ be an allowable digit set for β . If $T : \left[0, \frac{a_m}{\beta - 1}\right] \rightarrow \left[0, \frac{a_m}{\beta - 1}\right]$ is the greedy transformation with deleted digits for this β and A , then there exists a unique absolutely continuous, invariant measure, that is ergodic. Furthermore, the support of the probability density function f is the interval $[0, y_{i_0}]$, where $i_0 = \min\{i : T[0, y_i] \subseteq [0, y_i] \lambda \text{ a.e.}\}$.*

Now consider the lazy transformation with deleted digits for $\beta > 1$ and allowable digit set $\tilde{A} = \{\tilde{a}_0, \dots, \tilde{a}_m\}$, where $\tilde{a}_0 = 0$. Indicate the points of discontinuity of L in the following way. For $i \in \{0, \dots, m - 1\}$, let $\tilde{\ell}_i = \frac{\tilde{a}_m}{\beta - 1} - \frac{\tilde{a}_m - \tilde{a}_i}{\beta}$. In the same way as was done for the greedy transformation with deleted digits, we can make a partition $\{\tilde{\Delta}_i\}_{i=0}^m$, using these points of discontinuity. Let $\tilde{\Delta}_0 = [0, \tilde{\ell}_0]$ and $\tilde{\Delta}_m = \left(\tilde{\ell}_{m-1}, \frac{\tilde{a}_m}{\beta - 1}\right]$ and for all $i \in \{1, \dots, m - 1\}$, define

$$\tilde{\Delta}_i = (\tilde{\ell}_{i-1}, \tilde{\ell}_i].$$

For each $i \in \{1, \dots, m\}$, let \tilde{y}_i denote the value of L when taking the limit from the right to the point $\tilde{\ell}_{i-1}$, i.e. $\tilde{y}_i = \frac{\tilde{a}_m}{\beta - 1} - (\tilde{a}_i - \tilde{a}_{i-1})$. (See Figure 2)

The following corollary follows directly from Theorem 2.1.

Corollary 2.1. *Let L be the lazy transformation with deleted digits for $\beta > 1$ and allowable digit set $\tilde{A} = \{\tilde{a}_0, \dots, \tilde{a}_m\}$, for which $\tilde{a}_0 = 0$. Let T be the greedy transformation with deleted digits for the same $\beta > 1$ and allowable digit set $A = \{a_0, \dots, a_m\}$, such that $a_i = \tilde{a}_m - \tilde{a}_{m-i}$. Then there exists a unique absolutely continuous, invariant measure ν for L , that is ergodic. Let i_0 be defined for the greedy transformation as in equation (4). Then the support of the measure ν is given by the interval $\left[\tilde{y}_{m-i_0+1}, \frac{\tilde{a}_m}{\beta - 1}\right]$.*

Proof. By Theorem 2.1 we know that the interval $[0, y_{i_0}]$ is the support of the density function for the greedy transformation. Let μ be the absolutely continuous, invariant measure for T . Since T and L are isomorphic with isomorphism ϕ , given by $\phi(x) = \frac{a_m}{\beta - 1} - x$, L also has a unique absolutely continuous, invariant measure, given by $\nu = \mu \circ \phi^{-1}$. This is an ergodic measure and its support is given by

$$\phi([0, y_{i_0}]) = \left[\tilde{y}_{m-i_0+1}, \frac{\tilde{a}_m}{\beta - 1}\right]. \quad \square$$

3. EXAMPLES OF EXPLICITLY CALCULABLE INVARIANT MEASURES

In the previous section it is shown that by the results of T.Y. Li and J. Yorke in [8], T has a unique invariant measure that is absolutely continuous with respect to




FIGURE 2. The lazy β -transformation with $\beta = 2.5$ and $A = \{0, 1.35, 1.75, 3.3, 6\}$.

the normalized Lebesgue measure on $[0, y_{i_0}]$. The same observations can be made for the lazy transformation. In general we cannot give an explicit expression of this invariant measure, but below we discuss two cases of which we do know what the invariant measure is.

3.1. Imposing a condition on the number of digits. The first example is a particular case of the transformations that are studied by K. Wilkinson in [14]. In this paper, K. Wilkinson derives a formula for the density of an absolutely continuous, invariant measure for certain piecewise linear transformations. Before we go into more detail, let us give some definitions. We consider our greedy transformation T with $\beta > 1$ and $A = \{0, a_1, \dots, a_m\}$ an allowable digit set for β , but with the extra restriction that $m < \beta \leq m + 1$. In [4] it was shown that condition (1) implies that $\lceil \beta \rceil \leq m + 1$, so this example is the case in which we take as few digits as possible.

By the previous section, we know that the attractor is given by $[0, y_{i_0}]$. Suppose that N is the largest index such that $\frac{a_N}{\beta} < y_{i_0}$. Then the points $\frac{a_i}{\beta}$, $i = 1, \dots, N$, give an interval partition of the interval $[0, y_{i_0}]$ as before. Let $\Delta = \{\Delta_0, \dots, \Delta_N\}$ be the partition of $[0, y_{i_0}]$, such that

$$\Delta_0 = \left[0, \frac{a_1}{\beta}\right), \quad \Delta_N = \left[\frac{a_N}{\beta}, y_{i_0}\right]$$

and for $i = 1, \dots, N - 1$,

$$\Delta_i = \left[\frac{a_i}{\beta}, \frac{a_{i+1}}{\beta} \right).$$

An element $\Delta_i \in \Delta$ is called a *full interval of rank 1* if $\lambda(T\Delta_i) = 1$, where λ is the normalized Lebesgue measure on the interval $[0, y_{i_0}]$. Otherwise we call it *non-full*. Using Δ and T , we can make the sequence of partitions $\{\Delta^{(n)}\}$ in the usual way.

For $n \geq 1$, $\Delta^{(n)} = \bigvee_{k=0}^{n-1} T^{-k}\Delta$. The elements of $\Delta^{(n)}$ are intervals and are called

the *cylinder sets of order n* . An element $E^{(n)} \in \Delta^{(n)}$ is called *full of rank n* if $\lambda(T^n E^{(n)}) = 1$ and *non-full* otherwise. Now let $I(E^{(n)})$ be the number of non-full cylinders of rank $n + 1$, that are contained in $E^{(n)}$ and let

$$I_n = \sup_{E^{(n)} \in \Delta^{(n)}} I(E^{(n)}).$$

So for each cylinder set of rank n , we take the number of non-full subcylinder sets of rank $n + 1$ and I_n indicates the supremum of these numbers over all the cylinder sets of rank n . If we then take the supremum over all ranks, we get a number I , i.e.

$$I = \sup_{n \geq 0} I_n,$$

where I_0 is the number of non-full intervals of rank 1. In [14] K. Wilkinson derives a formula for the absolutely continuous invariant measure under the condition that $\beta > I$. We will adapt his result to our case and generalize it to our setting. For each $K \geq 0$, let $\bar{I}_K = \sup_{n \geq K} I_n$ and let B_n denote the union of those cylinder sets of order n which are full, but which are not a subset of any full cylinder set of lower rank. Notice that $I = \bar{I}_0$. We have the following lemma, which is a generalization of Corollary 4.5 in [14].

Lemma 3.1. *Let \bar{I}_K and B_n be as above and suppose that $\beta > \bar{I}_K$ for some $K \geq 0$. Then*

$$\sum_{n=1}^{\infty} \lambda(B_n) = 1.$$

Proof. Consider the attractor $[0, y_{i_0}]$ and cover it as far as possible with full intervals of rank 1. Since every non-full interval of rank 1 has Lebesgue measure smaller than $\frac{1}{\beta}$, the remaining part has Lebesgue measure smaller than $\frac{I_0}{\beta}$. Now fill this part as far as possible with full intervals of rank 2. The remaining part has Lebesgue measure smaller than $\frac{I_0 \cdot I_1}{\beta^2}$. If we continue in this manner, after $n + 1$ steps the remaining part will have Lebesgue measure smaller than

$$I_0 \cdot I_1 \cdot \dots \cdot I_n \cdot \frac{1}{\beta^n}.$$

And by hypothesis we have

$$\lim_{n \rightarrow \infty} I_0 \cdot I_1 \cdot \dots \cdot I_n \cdot \frac{1}{\beta^n} \leq I_0 \cdot I_1 \cdot \dots \cdot I_{K-1} \lim_{n \rightarrow \infty} \left(\frac{\bar{I}_K}{\beta} \right)^n = 0,$$

which completes the proof. \square

The next theorem is an adaptation of the formula by K. Wilkinson and a generalization of Theorem 5.12 of [14]. It gives an explicit expression of the absolutely continuous, invariant measure of the greedy transformation with deleted digits under the assumption that $\beta > \bar{I}_K$ for some $K \geq 0$. Before we state the theorem, we need the following notation. Let D_n be the union of all non-full cylinder sets of rank n , that are not subsets of any full cylinder set of lower rank. Let $x \in [0, y_{i_0}]$. Define $\phi_0(x) = 1$ and for $n \geq 1$, let

$$(5) \quad \phi_n(x) = \sum_{E^{(n)} \in D_n} \frac{1}{\beta^n} 1_{T^n E^{(n)}}(x).$$

Theorem 3.1. *If $\beta > \bar{I}_K$ for some $K \geq 0$, then the functions ϕ_n , $n \geq 0$ and ϕ , given by*

$$\phi : [0, y_{i_0}] \rightarrow [0, y_{i_0}] : x \mapsto \sum_{n=0}^{\infty} \phi_n(x)$$

are Lebesgue integrable and the function h given by

$$h : [0, y_{i_0}] \rightarrow [0, y_{i_0}] : x \mapsto \frac{\phi(x)}{\int \phi(x) d\lambda(x)}$$

is the absolutely continuous, invariant measure of T .

Proof. The proof follows from Lemma 3.1 and a slight adaptation of the corresponding proof in [14]. \square

Remark 3.1. In the case $K = 0$ Theorem 3.1 reduces to the theorem proved by K. Wilkinson.

The next theorem states the claim that in the case $m < \beta \leq m+1$, we have that $\beta > I$, so we can immediately apply Theorem 3.1.

Theorem 3.2. *Let $\beta > 1$ and $A = \{0, a_1, \dots, a_m\}$ be an allowable digit set, such that $m < \beta \leq m+1$. Let T be the greedy transformation for this β and A . Then the unique absolutely continuous, invariant density is given by Theorem 3.1.*

Proof. It is enough to show that $\beta > I$. First notice that Δ_{i_0-1} is a full interval, so that we have $I_0 \leq N \leq m < \beta$. By the definition of y_{i_0} we have that $\frac{a_{i_0}}{\beta} < y_{i_0}$, which means that $\Delta_{i_0-1} \neq \Delta_N$. Since for each cylinder set of order n , $E^{(n)}$, we have that $T^n E^{(n)}$ is an interval of the form $[0, y] \subseteq [0, y_{i_0}]$, we know that $E^{(n)}$ can contain at most N non-full intervals of rank n . So $I_n \leq N$ for each $n \geq 1$, which means that $I < \beta$, as we wanted. \square

We consider two examples. The first one satisfies the condition of Theorem 3.2. The second one does not satisfy the condition of this theorem, but in this case we can apply Theorem 3.1.

Example 3.1. First, let $\beta = 1 + \sqrt{2}$ be the positive solution of the equation $\beta^2 - 2\beta - 1 = 0$ and consider the allowable digit set $A = \{0, 1, 3\}$. The interval $[0, 2]$ is the attractor and Figure 3 gives the situation. The orbits of the points 1 and 2 are as follows.

$$\begin{aligned} T(1) &= \beta - 1, & T^2(1) &= T(\beta - 1) = \frac{1}{\beta}, & T^3(1) &= T\left(\frac{1}{\beta}\right) = 0, \\ T(2) &= 2\beta - 3, & T^2(2) &= T(2\beta - 3) = \beta - 1, & T^3(2) &= T(\beta - 1) = \frac{1}{\beta}. \end{aligned}$$




FIGURE 3. The greedy β -transformation with $\beta = 1 + \sqrt{2}$ and $A = \{0, 1, 3\}$ on the interval $[0, 2]$.

Notice that the condition of Theorem 3.1 is satisfied, so equation (5) gives for $x \in [0, 2]$,

$$\begin{aligned} \phi(x) &= 1 + \frac{1}{\beta} 1_{[0,1)}(x) + \frac{1}{\beta} 1_{[0,2\beta-3)}(x) + \sum_{k=0}^{\infty} \frac{c_{k+2}}{\beta^{k+2}} 1_{[0,1)}(x) \\ &\quad + \sum_{k=0}^{\infty} \frac{c_{k+1}}{\beta^{k+2}} 1_{[0,\beta-1)}(x) + \sum_{k=0}^{\infty} \frac{c_k}{\beta^{k+2}} 1_{[0,\frac{1}{\beta})}(x), \end{aligned}$$

where c_k is the k -th term of the tribonacci sequence, i.e. $c_k = c_{k-1} + c_{k-2} + c_{k-3}$, starting with $c_0 = 0$, $c_1 = c_2 = 2$. One can prove the following identity for the generating function of this sequence.

$$\sum_{k=0}^{\infty} c_k x^k = \frac{2x}{1 - x - x^2 - x^3}.$$

Using this formula, we get that

$$\begin{aligned} \sum_{k=0}^{\infty} \frac{c_{k+2}}{\beta^{k+2}} &= \frac{\frac{2}{\beta}}{1 - \frac{1}{\beta} - \frac{1}{\beta^2} - \frac{1}{\beta^3}} - \frac{2}{\beta} = 2 - \frac{1}{\beta}, \\ \sum_{k=0}^{\infty} \frac{c_{k+1}}{\beta^{k+2}} &= \frac{1}{\beta} \left[\frac{\frac{2}{\beta}}{1 - \frac{1}{\beta} - \frac{1}{\beta^2} - \frac{1}{\beta^3}} \right] = 1, \\ \sum_{k=0}^{\infty} \frac{c_k}{\beta^{k+2}} &= \frac{1}{\beta^2} \left[\frac{\frac{2}{\beta}}{1 - \frac{1}{\beta} - \frac{1}{\beta^2} - \frac{1}{\beta^3}} \right] = \frac{1}{\beta}. \end{aligned}$$

So, now

$$\phi(x) = 1 + \frac{1}{\beta} \cdot 1_{[0, 2\beta-3)}(x) + 2 \cdot 1_{[0, 1)}(x) + 1_{[0, \beta-1)}(x) + \frac{1}{\beta} \cdot 1_{[0, \frac{1}{\beta})}(x)$$

and rewriting this, leads to

$$\begin{aligned} \phi(x) = & 2\beta \cdot 1_{[0, \frac{1}{\beta})}(x) + (\beta + 2) \cdot 1_{[\frac{1}{\beta}, 1)}(x) + \beta \cdot 1_{[1, \beta-1)}(x) \\ & + (\beta - 1) \cdot 1_{[\beta-1, 2\beta-3)}(x) + 1_{[2\beta-3, 2)}(x). \end{aligned}$$

It can easily be verified that this is indeed an invariant density. Since

$$\int_{[0, 2)} \phi(x) d\lambda(x) = \frac{8\beta - 4}{\beta},$$

we know that the measure μ , given by

$$\mu(E) = \int_E \frac{\beta}{8\beta - 4} \phi(x) d\lambda(x),$$

for every measurable set E , is the unique invariant measure that is absolutely continuous with respect to the Lebesgue measure.

Example 3.2. For our second example we let $\beta = \frac{1 + \sqrt{5}}{2}$ be the golden mean, i.e. the positive solution of the equation $\beta^2 - \beta - 1 = 0$, and we take as our allowable digit set $A = \{0, 2\beta, 5\}$. Notice that with this combination of β and A the condition of Theorem 3.2 is not satisfied. The attractor is given by the interval $[0, 2\beta]$ and in Figure 4 the transformation T is given on this interval. We now look at the orbits




FIGURE 4. The greedy β -transformation with $\beta = \frac{1 + \sqrt{5}}{2}$ and $A = \{0, 2\beta, 5\}$ on the interval $[0, 2\beta]$.

of the points 2β and $5 - 2\beta$.

$$\begin{aligned} T(2\beta) &= 2\beta - 3, & T^2(2\beta) &= 2 - \beta, & T^3(2\beta) &= \beta - 1, & T^4(2\beta) &= 1, \\ T^5(2\beta) &= \beta + 1, & T^6(2\beta) &= 1, \end{aligned}$$

$$\begin{aligned} T(5 - 2\beta) &= 3\beta - 2, & T^2(5 - 2\beta) &= 3 - \beta, & T^3(5 - 2\beta) &= 2\beta - 1, \\ T^4(5 - 2\beta) &= 2 - \beta. \end{aligned}$$

And we see that after the first iteration both orbits never hit Δ_2 again. This means that the number of non-full intervals of rank $n \geq 2$ that are not a subset of any full interval of lower rank is at most 1, hence $\bar{I}_2 = 1 < \beta$. By Theorem 3.1 we can apply formula (5) to get

$$\begin{aligned} \phi(x) &= 1 + \frac{1}{\beta} \cdot 1_{[0,5-2\beta)}(x) + \frac{1}{\beta} \cdot 1_{[0,2\beta-3)}(x) + \frac{1}{\beta^2} \cdot 1_{[0,3\beta-2)}(x) + \frac{2}{\beta^3} \cdot 1_{[0,2-\beta)}(x) \\ &\quad + \frac{1}{\beta^3} \cdot 1_{[0,3-\beta)}(x) + \frac{2}{\beta^4} \cdot 1_{[0,\beta-1)}(x) + \frac{1}{\beta^4} \cdot 1_{[0,2\beta-1)}(x) \\ &\quad + \left[\frac{1}{\beta^4} + \frac{2}{\beta^7} \sum_{k=0}^{\infty} \frac{1}{\beta^{3k}} \right] 1_{[0,1)}(x) + \left[\frac{1}{\beta^5} + \frac{2}{\beta^8} \sum_{k=0}^{\infty} \frac{1}{\beta^{3k}} \right] 1_{[0,\beta)}(x) \\ &\quad + \left[\frac{1}{\beta^6} + \frac{2}{\beta^9} \sum_{k=0}^{\infty} \frac{1}{\beta^{3k}} \right] 1_{[0,\beta+1)}(x). \end{aligned}$$

Rewriting this will get you

$$\begin{aligned} \phi(x) &= (2\beta + 1) \cdot 1_{[0,2\beta-3)}(x) + (\beta + 2) \cdot 1_{[2\beta-1,2-\beta)}(x) + (8 - 3\beta) \cdot 1_{[2-\beta,\beta-1)}(x) \\ &\quad + (3\beta - 2) \cdot 1_{[\beta-1,1)}(x) + (\beta + 1) \cdot 1_{[1,3-\beta)}(x) + (4 - \beta) \cdot 1_{[3-\beta,\beta)}(x) \\ &\quad + (2\beta - 1) \cdot 1_{[\beta,5-2\beta)}(x) + \beta \cdot 1_{[5-2\beta,2\beta-1)}(x) + (4\beta - 5) \cdot 1_{[2\beta-1,\beta+1)}(x) \\ &\quad + (3 - \beta) \cdot 1_{[\beta+1,3\beta-2)}(x) + 1_{[3\beta-2,2\beta)}(x). \end{aligned}$$

Furthermore, we have

$$\int_0^{2\beta} \phi(x) d\lambda(x) = \frac{49 - 23\beta}{\beta},$$

so the density h of the unique absolutely continuous, invariant measure is given by

$$h(x) = \frac{\beta\phi(x)}{49 - 23\beta}.$$

Remark 3.2. In [14] more is said about piecewise linear transformations with maximal slope β for which $I < \beta$. For example, K. Wilkinson proves that these transformations are exact and weak Bernoulli. We can remark that greedy β -transformations with deleted digit for which the number of digits $m + 1$ satisfies $m < \beta \leq m + 1$ has these same properties, i.e. they are exact and weak Bernoulli.

3.2. Ultimately periodic endpoints. The condition we impose on the system in this second example is that the endpoints of the transformation must have ultimately periodic orbits. What we mean by this is clarified in the following definition. Let T be the greedy transformation with deleted digits for $\beta > 1$ and allowable digit set $A = \{0, a_1, \dots, a_m\}$, considered on its attractor $[0, y_{i_0}]$ as given in Theorem 2.1. Let $N \geq 1$ be the largest index such that $\frac{a_N}{\beta} < y_{i_0}$. We say that *the endpoints of T have ultimately periodic orbits* if for each $i \in \{1, \dots, N\}$ there exist numbers $u(i), p(i)$, such that $T^{u(i)+p(i)}y_i = T^{u(i)}y_i$. In this case we say that the points y_i have ultimately periodic orbits of period $p(i)$.

In [2] W. Byers and A. Boyarsky proved some nice results about the absolutely continuous, invariant measure of a certain class of piecewise linear functions, namely the piecewise linear Markov maps, which they defined as follows. Let $0 = \alpha_0 < \alpha_1 < \dots < \alpha_n = 1$ be a partition of the interval $[0, 1]$, denoted by \mathcal{P} . A map $\tau : [0, 1] \rightarrow [0, 1]$ is called a *piecewise linear Markov map* if $\tau|_{(\alpha_i, \alpha_{i+1})}$ is a linear homeomorphism onto some interval $(\alpha_{j(i)}, \alpha_{k(i)})$ for all $i \in \{0, \dots, n-1\}$. If τ is such a map, it induces an $n \times n$ 0-1 matrix $M = M_\tau$ in the following way. The entry m_{ij} equals 1 if $[\alpha_j, \alpha_{j+1}] \subseteq \tau([\alpha_i, \alpha_{i+1}])$ and 0 otherwise. The fact that τ is a piecewise linear Markov map guarantees that the nonzero entries in each row are contiguous. In [2], W. Byers and A. Boyarsky proved the following results.

Let τ be a piecewise linear Markov map, that is expanding and of constant slope and suppose that the 0-1 matrix M it induces is irreducible. Then

(d1) There exists a unique invariant probability measure μ , that is equivalent to the Lebesgue measure and that maximizes entropy.

(d2) The entropy of τ with respect to this measure μ equals $\log \beta$, where β is the spectral radius of M and is also equal to the slope of τ .

A combination of these results, with those found by W. Parry in [10], gives that the invariant measure μ of τ can be found in the following way. Let β be as given by (d2). Let $\mathbf{v} = (v_0, \dots, v_n)$ be the right eigenvector of M , belonging to the eigenvalue β and such that $\sum_{i=0}^n v_i = 1$ and suppose that $\mathbf{u} = (u_0, \dots, u_n)$ is the left

eigenvector of M belonging to eigenvalue β and such that $\sum_{i=0}^n u_i v_i = 1$. Then the function

$$(6) \quad \phi : [0, 1] \rightarrow [0, 1] : x \mapsto \sum_{i=0}^n u_i \cdot 1_{[\alpha_i, \alpha_{i+1})}(x)$$

is the density of the unique, absolutely continuous, invariant measure for τ that we are looking for.

Now, consider the greedy transformation with deleted digits T that has ultimately periodic endpoints. Using the orbits of these endpoints, we make a partition \mathcal{P} of the interval $[0, y_{i_0}]$. Consider the set

$$I = \{T^k y_i : 1 \leq i \leq N, k \geq 0\} \cup \left\{ \frac{a_i}{\beta} : 1 \leq i \leq N \right\} \cup \{0\}.$$

Since all the orbits of the points y_i are periodic, this set only contains a finite number of elements, say $n+1$ elements, so we can put them in increasing order, to get a sequence $0 = \alpha_0 < \alpha_1 < \dots < \alpha_n = y_{i_0}$. This gives us the partition \mathcal{P} , i.e. $\mathcal{P} = \{P_i\}_{i=0}^n$ with $P_i = [\alpha_i, \alpha_{i+1})$. The next lemma states that in this case T is a piecewise linear Markov map for this partition.

Lemma 3.2. *Let T and \mathcal{P} be as above. Then T is linear on each of the elements of \mathcal{P} and for each $i \in \{0, \dots, n\}$ we have $TP_i = [\alpha_{j(i)}, \alpha_{k(i)})$, for some $\alpha_{j(i)}, \alpha_{k(i)} \in I$.*

Proof. Since all the points $\frac{a_i}{\beta}$ are in I , it is easy to see that T is linear on each element of \mathcal{P} . Now fix an element $P_i \in \mathcal{P}$. If $\alpha_i = \frac{a_j}{\beta}$ for some $j \in \{1, \dots, N\}$,

then $TP_i = [0, T^k y_\ell)$ for some $\ell \in \{1, \dots, N\}$ and $k \geq 0$. Suppose $\alpha_i = T^k y_\ell$. If $\alpha_{i+1} = \frac{\alpha_j}{\beta}$, then $T\alpha_i < y_j$, so $TP_i = [T^{k+1} y_\ell, y_j)$. If, on the other hand $\alpha_{i+1} = T^m y_j$, then $T\alpha_i < T\alpha_{i+1}$ and $TP_i = [T^{k+1} y_\ell, T^{m+1} y_j)$. In all cases TP_i is of the desired form. \square

The following lemma states that each element of \mathcal{P} is eventually mapped in each other element of \mathcal{P} , which means that the matrix that T induces is irreducible.

Lemma 3.3. *For each $i, j \in \{0, \dots, n\}$ there exists an $k \geq 0$, such that $P_j \subseteq T^k P_i$.*

Proof. Let P_i and P_j be two elements of \mathcal{P} . Let μ be the measure given by Theorem 2.1 and let ϕ be its density. By (c4) we know that $\phi > 0$ almost everywhere on P_i and P_j , so P_i and P_j both have strictly positive measure. Moreover, μ is ergodic, so there exists a $k \geq 0$, such that

$$(7) \quad \mu(P_i \cap T^{-k} P_j) > 0.$$

By Lemma 3.2, we have that $T^k P_i = \bigcup_{\ell \in \alpha} P_\ell$ for some index set $\alpha \subseteq \{0, \dots, n\}$.

Then (7) gives that $\mu(P_j \cap \bigcup_{\ell \in \alpha} P_\ell) > 0$, so that $P_j \subseteq T^k P_i$. \square

It is easy to see that the transformation T , defined on its attractor $[0, y_{i_0}]$ is isomorphic to a greedy transformation with deleted digits \bar{T} , defined on the interval $[0, 1]$. The isomorphism ψ is given by $\psi : [0, 1] \rightarrow [0, y_{i_0}] : x \mapsto y_{i_0} x$ and we have $T \circ \psi = \psi \circ \bar{T}$. The transformation T induces an $n \times n$ 0-1 matrix M in the following way. The entry $m_{ij} = 1$ if $P_j \subseteq TP_i$ and $m_{ij} = 0$ otherwise. By Lemma 3.2 the ones in each row of M are consecutive and by Lemma 3.3, the matrix M is irreducible. Let $\mathbf{v} = (v_0, \dots, v_n)$ be the right eigenvector of M , belonging to the eigenvalue β and such that $\sum_{i=0}^n v_i = 1$ and suppose that $\mathbf{u} = (u_0, \dots, u_n)$ is the left eigenvector

of M belonging to eigenvalue β and such that $\sum_{i=0}^n u_i v_i = 1$. The transformation \bar{T} induces the same matrix and therefore, we have the following theorem.

Theorem 3.3. *Let $\beta > 1$ and $A = \{0, a_1, \dots, a_m\}$ be an allowable digit set. Consider T on the attractor $[0, y_{i_0}]$ and suppose that all the endpoints of T have periodic orbits. Let \bar{T} be the greedy transformation with deleted digits, that is defined on $[0, 1]$ and that is isomorphic to T by the isomorphism ψ defined above. Then the unique absolutely continuous, invariant measure μ_T is the unique measure that maximizes entropy. This entropy is given by $\log \beta$. The measure μ_T is defined for all measurable sets B by*

$$\mu_T(B) = \mu_{\bar{T}}(\psi^{-1}(B)),$$

where $\mu_{\bar{T}}$ is the absolutely continuous, invariant measure for \bar{T} , of which the probability density function is given by equation (6).

We consider the same two examples as in the previous paragraph.

Example 3.3. In the first example, $\beta = 1 + \sqrt{2}$ was the positive solution of the equation $\beta^2 - 2\beta - 1 = 0$ and we considered the allowable digit set $A = \{0, 1, 3\}$.

The attractor was the interval $[0, 2]$. We already saw that both endpoints have finite orbits and the partition $\mathcal{P} = \{P_i\}_{i=0}^5$, given by these orbits is as follows:

$$\begin{aligned} P_0 &= \left[0, \frac{1}{\beta}\right), & P_1 &= \left[\frac{1}{\beta}, 1\right), & P_2 &= \left[1, \frac{3}{\beta}\right), \\ P_3 &= \left[\frac{3}{\beta}, \beta - 1\right), & P_4 &= [\beta - 1, 2\beta - 3], & P_5 &= [2\beta - 3, 2]. \end{aligned}$$

This gives us the following 0-1 matrix:

$$M = \begin{pmatrix} 1 & 1 & 0 & 0 & 0 & 0 \\ 1 & 1 & 1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 1 \\ 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1 & 1 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 \end{pmatrix}.$$

Let $\mathbf{v} = (v_0, \dots, v_5)$ be the right eigenvector of M with eigenvalue β and such that $\sum_{i=0}^5 v_i = 1$ and let $\mathbf{u} = (u_0, \dots, u_5)$ be the left eigenvector of M for the eigenvalue β and such that $\sum_{i=0}^5 u_i v_i = 1$. Then

$$\mathbf{v} = \frac{1}{2\beta^2}(\beta, \beta + 1, \beta - 1, 1, \beta, 1)$$

and

$$\mathbf{u} = \frac{\beta}{4\beta - 2}(2\beta, \beta + 2, \beta, \beta, \beta - 1, 1).$$

This means that the invariant probability density for our transformation T is given by

$$\begin{aligned} h(x) &= \frac{1}{2} \cdot \frac{\beta}{4\beta - 2} [2\beta \cdot 1_{[0, \frac{1}{\beta})}(x) + (\beta + 2) \cdot 1_{[\frac{1}{\beta}, 1)}(x) + \beta \cdot 1_{[1, \beta - 1)}(x) \\ &\quad + (\beta - 1) \cdot 1_{[\beta - 1, 2\beta - 3)}(x) + 1_{[2\beta - 3, 2)}(x)], \end{aligned}$$

just as we obtained before.

Example 3.4. In the second example, $\beta = \frac{1 + \sqrt{5}}{2}$ was the golden mean and $A = \{0, 2\beta, 5\}$ was the allowable digit set. The attractor was $[0, 2\beta]$. In this case, both the orbit of 2β and that of $5 - 2\beta$ were eventually periodic and the partition $\mathcal{P} = \{P_i\}_{i=0}^{12}$ and 0-1 matrix M they give are the following. For the partition we have

$$\begin{aligned} P_0 &= [0, 2\beta - 3), & P_1 &= [2\beta - 3, 2 - \beta), & P_2 &= \left[2 - \beta, \frac{1}{\beta}\right), \\ P_3 &= \left[\frac{1}{\beta}, 1\right), & P_4 &= [1, 3 - \beta), & P_5 &= [3 - \beta, \beta), \\ P_6 &= [\beta, 5 - 2\beta), & P_7 &= [5 - 2\beta, 2), & P_8 &= [2, 2\beta - 1), \\ P_9 &= [2\beta - 1, \beta + 1), & P_{10} &= [\beta + 1, 3\beta - 2), & P_{11} &= \left[3\beta - 2, \frac{5}{\beta}\right), \\ P_{12} &= \left[\frac{5}{\beta}, 2\beta\right]. \end{aligned}$$

And for the matrix,

$$M = \begin{pmatrix} 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 \\ 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 1 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \end{pmatrix}.$$

The right eigenvector, \mathbf{v} , with eigenvalue β and such that the sum of its elements equals 1 is

$$\mathbf{v} = \frac{1}{10\beta + 6}(\beta, 1, \beta, \beta + 1, \beta + 1, \beta, 1, \beta, \beta, \beta + 1, \beta, \beta, 1)$$

and the left eigenvector, \mathbf{u} , belonging to the eigenvalue β and such that the dot product with \mathbf{v} is 1 is

$$\mathbf{u} = \frac{10\beta + 6}{29\beta + 3}(2\beta + 1, 2 + \beta, 8 - 3\beta, 3\beta - 2, \beta + 1, 4 - \beta, 2\beta - 1, \beta, \beta, 4\beta - 5, 3 - \beta, 1, 1).$$

The invariant probability density then is

$$\begin{aligned} h(x) &= \frac{1}{2\beta} \cdot \frac{10\beta + 6}{29\beta + 3} [(2\beta + 1) \cdot 1_{[0, 2\beta - 3)}(x) + (\beta + 2) \cdot 1_{[2\beta - 1, 2 - \beta)}(x) \\ &\quad + (8 - 3\beta) \cdot 1_{[2 - \beta, \beta - 1)}(x) + (3\beta - 2) \cdot 1_{[\beta - 1, 1)}(x) + (\beta + 1) \cdot 1_{[1, 3 - \beta)}(x) \\ &\quad + (4 - \beta) \cdot 1_{[3 - \beta, \beta)}(x) + (2\beta - 1) \cdot 1_{[\beta, 5 - 2\beta)}(x) + \beta \cdot 1_{[5 - 2\beta, 2\beta - 1)}(x) \\ &\quad + (4\beta - 5) \cdot 1_{[2\beta - 1, \beta + 1)}(x) + (3 - \beta) \cdot 1_{[\beta + 1, 3\beta - 2)}(x) + 1_{[3\beta - 2, 2\beta)}(x)]. \end{aligned}$$

And again, this is equal to the result from the previous paragraph.

4. CONCLUSIONS

In the second section of this article we have established that the greedy β -transformation with deleted digits has a unique invariant measure that is absolutely continuous with respect to the Lebesgue measure. We saw that this measure is ergodic and gave the interval on which the density function is strictly positive.

In the last section we have studied two specific examples of the greedy transformation in which this absolutely continuous, invariant measure can be explicitly calculated. The first example put a restraint on the number of digits that can be chosen. If this number $m + 1$, satisfies $m < \beta \leq m + 1$, then the density of the absolutely continuous, invariant measure is given by K. Wilkinson's formula. We remarked that in this case the absolutely continuous, invariant measure is exact and weak Bernoulli. The second example we studied was the case in which the endpoints of the transformation have ultimately periodic orbits. In that case the absolutely continuous, invariant measure is also the measure of maximal entropy (with entropy equal to $\log \beta$) and its density is given by W. Parry's formula.

REFERENCES

- [1] J. Buzzi and O. Sarig. Uniqueness of equilibrium measures for countable Markov shifts and multidimensional piecewise expanding maps. *Ergodic Theory Dynam. Systems*, 23(5):1383–1400, 2003.
- [2] W. Byers and A. Boyarsky. Absolutely continuous invariant measures that are maximal. *Trans. Amer. Math. Soc.*, 290(1):303–314, 1985.
- [3] W. Byers, P. Góra, and A. Boyarsky. Maximal absolutely continuous invariant measures for piecewise linear Markov transformations. *Ergodic Theory Dynam. Systems*, 10(4):645–656, 1990.
- [4] K. Dajani and C. Kalle. Random β -expansions with deleted digits. *To appear in Disc. and Cont. Dyn. Sys.* Preprint available at <http://www.math.uu.nl/publications/preprints/1354.pdf>.
- [5] F. Hofbauer. Maximal measures for piecewise monotonically increasing transformations on $[0, 1]$. In *Ergodic theory (Proc. Conf., Math. Forschungsinst., Oberwolfach, 1978)*, volume 729 of *Lecture Notes in Math.*, pages 66–77. Springer, Berlin, 1979.
- [6] F. Hofbauer. On intrinsic ergodicity of piecewise monotonic transformations with positive entropy. *Israel J. Math.*, 34(3):213–237 (1980), 1979.
- [7] A. Lasota and J. A. Yorke. On the existence of invariant measures for piecewise monotonic transformations. *Trans. Amer. Math. Soc.*, 186:481–488 (1974), 1973.
- [8] T. Y. Li and J. A. Yorke. Ergodic transformations from an interval into itself. *Trans. Amer. Math. Soc.*, 235:183–192, 1978.
- [9] W. Parry. On the β -expansions of real numbers. *Acta Math. Acad. Sci. Hungar.*, 11:401–416, 1960.
- [10] W. Parry. Intrinsic Markov chains. *Trans. Amer. Math. Soc.*, 112:55–66, 1964.
- [11] M. Pedicini. Greedy expansions and sets with deleted digits. *Theoret. Comput. Sci.*, 332(1-3):313–336, 2005.
- [12] A. Rényi. Representations for real numbers and their ergodic properties. *Acta Math. Acad. Sci. Hungar.*, 8:477–493, 1957.
- [13] Fritz Schweiger. Some remarks on ergodicity and invariant measures. *Michigan Math. J.*, 22(2):181–187, 1975.
- [14] K. Wilkinson. Ergodic properties of a class of piecewise linear transformations. *Z. Wahrscheinlichkeitstheorie und Verw. Gebiete*, 31:303–328, 1974/75.

FAC. WISKUNDE EN INFORMATICA, UNIVERSITEIT UTRECHT, POSTBUS 80.000, 3508 TA UTRECHT,
THE NETHERLANDS

E-mail address: `dajani@math.uu.nl`

FAC. WISKUNDE EN INFORMATICA, UNIVERSITEIT UTRECHT, POSTBUS 80.000, 3508 TA UTRECHT,
THE NETHERLANDS

E-mail address: `kalle@math.uu.nl`